



Research Article

Spectrum of genetic variation in different selection schemes of desi chickpea (*Cicer arietinum* L.)

M. R. Paneliya, D. R. Mehta, J. Raval Lata and C. P. Chetariya

Department of Genetics and Plant Breeding, Junagadh Agricultural University, Junagadh-36200, Gujarat.

E-Mail: paneliyamehul.mp@gmail.com

(Received:07 Oct 2017; Revised:26 Aug 2018; Accepted: 26 Aug 2018)

Abstract

An experiment was conducted to study the spectrum of genetic variation for seed yield per plant and its component traits in the four selection procedures [PS(EF), PS(HY), SSD and RBP] each with 20 progenies of GJG 0315 x ICCV 96029 cross in F₅ generation in Desi chickpea (*Cicer arietinum* L.). Analysis of variance revealed significant genotypic differences for all the characters with a wide range of variation. Out of the total 6 cases studied in F₅, PS(EF) in 2 cases; SSD in 3 cases and PS(HY) in 1 cases were better for depicting the widest phenotypic range and maximum coefficient of variation irrespective of characters. Moderate to high heritability and genotypic co-efficients of variation coupled with high expected Genetic Advance as per cent of mean was observed for number of branches per plant by PS(HY) and RBP and for 100-seed weight in PS(EF), PS(HY) and SSD of this cross which indicated the predominant role of additive gene action in the expression of these traits in respective breeding schemes.

Keywords

Breeding procedures, genetic variability, chickpea, Desi

Introduction

Pulses are the most important source of vegetarian protein, high in fiber content and provide ample quantity of vitamins and minerals. Keeping in view large benefits of pulses for human health, the United Nations has proclaimed 2016 as the International Year of Pulses (Sandhu, 2015). Chickpea (*Cicer arietinum* L.), one of the major pulses cultivated and consumed in India, is also known as Bengal gram and is a major and cheap source of protein as compared to animal protein. In grain legumes, the improvement in seed yield through selection has not been encouraging due to its complex polygenic nature. Various breeding procedures have their own advantages and limitations. The variability of a biological population is an outcome of genetic constitution of the individuals making up of that population in relation to the prevailing environments. An assessment of genetic variation with the help of parameters such as genotypic coefficient of variation, heritability estimates and genetic advance are absolutely necessary to commence an efficient breeding programme. All the attempts about genetic improvement in seed yield and its economic return in chickpea are now directed towards the manipulation of genetic variation through hybridization followed by handling segregating generation with selection schemes.

Materials and Methods

The comparison of four selection procedures *viz.*,

pedigree selection for early flowering [PS(EF)], pedigree selection for high yield [PS(HY)], single seed descent (SSD) and random bulk population (RBP) were evaluated in F₅ generations of chickpea cross GJG 0315 x ICCV 96029. A total of 80 progenies (20 progenies in each selection scheme a cross) were evaluated in F₅ along with original F₂ and two parental lines during Rabi 2015-16 in Randomized Block Design (RBD) with three replications.

Initial crosses were made in Rabi 2010-11 at Pulse Research Station, JAU, Junagadh, F₁ in Rabi 2011-12, F₂ seeds from bulk of F₁ was raised in Rabi 2012-13 (75% used for selection and rest 25 % reserved for comparison in F₅), F₃ in Rabi 2013-14, F₄ in Rabi 2014-15 and F₅ in Rabi 2015-16 (25% saved F₂ was used.). Upto five years there was no harm to use the old seeds but viability was slightly declined. Observations were recorded on five randomly selected plants in each entry and replication for seed yield per plant and its component traits *viz.*, number of branches per plant, number of pods per plant, biological yield per plant, 100-seed weight and harvest index and their mean values were used for the statistical analysis. The genotypic (GCV) and phenotypic (PCV) coefficients of variation were estimated as per the formulae suggested by Burton (1952), while heritability in broad-sense and genetic advance were calculated by using the formulae suggested by Allard (1960).

Results and Discussion

The analysis of variance in F₅ generation indicated that all the four selection procedures ([PS(EF)], [PS(HY)], SSD and RBP), two parents and F₂ population differed significantly for all the characters indicating presence of sufficient amounts variability among all the four methods including two parents and F₂ population.

A wider phenotypic range and coefficient of variability was noted among progenies from PS(EF) compared to other selection procedures for number of pods per plant (41.33 – 65.33, 22.50%) and biological yield per plant (19.66 – 32.34, 24.38%). Similarly, wider range was also observed among lines derived from SSD for number of branches per plant (2.60 – 6.20, 40.91%), seed yield per plant (9.13 – 13.87, 20.61%) and harvest index (40.97 – 67.26, 24.29%) and PS(HY) for 100-seed weight (12.43 – 16.30, 13.47%). The variation in phenotypic range for different quantitative characters in F₄ derived lines in F₅ generation could be attributed to the substantial change brought about by selection in genetic makeup of chickpea through different selection procedures.

Among different selection procedures (Table 1), PS(HY) was numerically superior for biological yield per plant (27.12g). However, PS(HY) was found less effective to rest of the selection procedures for number of pod per plant (50.23g), 100-seed weight (14.59g) and seed yield per plant (10.02g). The effectiveness of early generation selection for seed yield was reported by Ivers and Fehr (1978) in soybean which contradictory to the present findings. PS(HY), however, did not turn out to be superior in this cross for seed yield per plant in the present study. Such observations were reported by Byth *et al.* (1979) in chickpea as well as in soybean by Pushpendra and Ram (1987). From time to time, several reasons have been proposed for failure of isolating high yielding plants in early segregating generations. With the very large genotypic variation available from F₂ populations, segregating generations no longer could be handled *via* pedigree selection.

Significantly better mean performance of SSD over other selection procedures was observed for 100-seed weight per plant (19.40g), seed yield per plant (11.00g) and harvest index (54.00%). For number of pods per plant, SSD (50.72) was found to be superior to PS(HY) (50.23). Thus, SSD seemed to be an effective alternative in case when it is not possible for a breeder to handle large segregating materials with limited resources. The SSD procedure has been shown often superior or at least equally efficient to traditional methods for developing superior lines by Sharma and Chaudhary (1989) in chickpea. RBP was found

superior for number of branches per plant (5.13) and number of pods per plant (52.95) and also RBP (15.02) was found superior to PS(HY) (14.59) for 100-seed weight. Under the circumstances, SSD and RBP methods have been found to be useful for carrying such populations (Frey, 1957). If direct selection was not effective in early generations, either SSD or RBP methods could be the best alternative for advancing populations in later generations.

Coefficient of variation measures the relative amounts of variation for different characters by bringing various measure of dispersion on a uniform scale and are, therefore, comparable. The better index for measuring the genetic variability is genotypic coefficient of variation (GCV%) as described by Burton (1952). The high values of GCV and PCV were observed with PS(EF) for number of pods per plant (10.11%, 13.69%) and seed yield per plant (11.04%, 16.62%); with SSD for harvest index (10.91%, 17.50%) and with RBP for number of branches per plant (20.57%, 24.15%) and biological yield per plant (9.18%, 13.32%), respectively. Irrespective of different selection methods, the cross exhibited high GCV (%) and PCV (%) for number of branches per plant. High PCV was observed for number of branches per plant in PS(HY) (20.67%), SSD (22.80%) and RBP (24.15%). This suggested that the greater variability for this character among lines had genetic basis and could be improved through selection. Selection based on phenotypic performance would be effective for improvement of seed yield and its component traits by different selection procedures. Such conclusions have been reached by Mehta and Zaveri, (1994).

In the present investigation, high heritability values was recorded for 100-seed weight in PS(EF) (61.09%), PS(HY) (87.80%) and SSD (94.98%). High heritability for 100-seed weight was also reported by Gul *et al.* (2013) and Monpara and Gaikwad (2014). Similarly high expected genetic advance (per cent of mean) was recorded for number of branches per plant in three selection procedures *viz.*, PS(HY) (30.07%), SSD (27.68%) and RBP (36.10%). High heritability (72.54%) along with high genetic advance (36.10%) was noted for number of branches per plant in RBP which indicated the role of additive genetic variance as reported by Kumar *et al.* (2012) and Neelu Kumari *et al.* (2013). High estimate of broad sense heritability was observed for most of the characters in different selection procedures of this cross. This indicated that the magnitude of heritability varies more as a function of the genetic variability and of the adaptive or constitutive nature of genetic differences, than as a function of the environment (Ceccarelli, 1989). He also suggested



that the magnitude of heritability does not necessarily represent the best criterion to use in deciding the optimal environment for selection.

Overall high heritability coupled with high genetic advance, expressed as per cent of mean, for most of the characters in different combinations of selection procedures indicated that these characters were under the control of additive gene action. Therefore, different selection procedures were effective in bringing desirable improvement in these characters.

It is concluded from the present study that among different selection procedures, high heritability along with moderate to high genetic advance, as per cent of mean, was observed for number of branches per plant in PS(HY) and RBP and for 100-seed weight in three selection schemes [PS(EF), PS(HY) and SSD] in this cross.

References

- Allard, R. W. 1960. Principles of Plant Breeding. John Wiley and Sons, New York.
- Burton, G. W. 1952. Quantitative inheritance in grasses. Proc. 6th Int. Grassland Cong., **1**: 277-283.
- Byth, D. E.; Green, J. M. and Hawtin, G. C. 1979. ICRISAT/ICARDA chickpea breeding strategies. In: Proceedings of the International Workshop on Chickpea Improvement. ICRISAT, Hyderabad. pp. 11-27, 28 Feb. to 2 March, 1979.
- Ceccarelli, S. 1989. Wide adaptation, How wide? *Euphytica*, **40**:197-205.
- Frey, K. J. and Horner, T. 1957. Heritability in standard units. *Agron. J.*, **49**:59-62.
- Gul, R.; Khan, H.; Bibi, M.; Ain, Q. U. and Imran, B. 2013. Genetic analysis and interrelationship of yield attributing traits in chickpea (*Cicer arietinum* L.). *The J. Animal & Plant Sci.*, **23**(2): 521-526.
- Ivers, F. H. and Fehr, W. R. 1978. Evaluation of the pureline family method for cultivar development. *Crop Sci.*, **18**: 541-544.
- Kumar, A.; Suresh Babu, G. and Roopa Lavanya, G. 2012. Character association and path analysis in early segregating population in chickpea (*Cicer arietinum* L.). *Legume Res.*, **35**: 337-340.
- Mehta, D. R. and Zaveri, P. P. 1994. Comparison of five selection schemes in four cowpea crosses. In: international symposium on pulse research, held during Agril. 2-6, 1994 at New Delhi.
- Monpara, B. A. and Gaikwad, S. R. 2014. Combining high seed number and weight to improve seed yield potential of chickpea in India. *African Crop Sci. J.*, **22**: 1-7.
- Neelu Kumari; Babu, S. and Lavanya, G. R. 2013. Genetic variability and character association in chickpea germplasm. *Trends in Bio. Sci.*, **6**:742-743.
- Pushpendra and Ram, H. H. 1987. Early generation selection for number of pods, harvest index and yield in soybean. *Crop Sci.*, **14**: 123-127.
- Sandhu, J. C. 2015. Status of pulse in India- India Pulses and Grain Association. (<http://www.ipga.co.in/status-of-pulses-in-india>).
- Sharma, R. N. and Chaudhary, J. S. 1989. Relative efficacy of single seed descent (SSD), pedigree selection (PS) and bulk population (BP) methods of breeding in chickpea. In: National Symposium on New Frontiers in Pulses Research and Development, Kanpur, November 10-12, 1989.



Table 1. Phenotypic range, mean and variability parameters for various traits in F₅ generation of desi chickpea

Selection procedure	Phenotypic range	Coefficient of variation (%)	Mean ± S.E.	PCV (%)	GCV (%)	h ² (%)	GA (% mean)
Number of branches per plant							
PS (EF)	3.60-5.93	24.45	4.44 ± 0.43	19.55	9.58	24.00	9.67
PS (HY)	3.00-6.40	36.17	4.63 ± 0.29	20.67	17.37	70.63	30.07
SSD	2.60-6.20	40.91	4.59 ± 0.38	22.80	17.50	58.94	27.68
RBP	3.00-7.00	40.00	5.13 ± 0.37	24.15	20.57	72.54	36.10
F ₂	2.00-3.67	29.45	2.73 ± 0.41	-	-	-	-
P ₁	3.33-7.00	35.33	5.07 ± 0.71	-	-	-	-
P ₂	2.33-5.67	41.75	3.73 ± 0.56	-	-	-	-
Number of pods per plant							
PS (EF)	41.33-65.33	22.50	51.81 ± 2.69	13.69	10.11	54.52	15.38
PS (HY)	43.33-57.33	13.91	50.23 ± 2.95	12.07	6.06	25.20	6.27
SSD	44.33-62.00	16.62	50.72 ± 2.78	13.18	8.89	45.37	12.32
RBP	46.33-68.67	19.43	52.95 ± 2.85	12.25	7.65	38.96	9.84
F ₂	31.00-95.00	50.79	51.87 ± 8.08	-	-	-	-
P ₁	42.67-57.33	14.66	50.73 ± 4.77	-	-	-	-
P ₂	42.33-53.33	11.50	48.47 ± 2.31	-	-	-	-
Biological yield per plant (g)							
PS (EF)	19.66-32.34	24.38	24.59 ± 1.31	13.00	8.92	47.10	12.62
PS (HY)	22.75-31.78	16.56	27.12 ± 1.41	12.26	8.06	43.22	10.91
SSD	19.29-25.54	13.94	22.46 ± 1.19	11.30	6.23	30.39	7.08
RBP	23.07-32.48	16.94	26.82 ± 1.46	13.32	9.18	47.51	13.04
F ₂	5.87-51.30	79.46	30.37 ± 3.53	-	-	-	-
P ₁	21.40-28.53	14.28	24.20 ± 1.70	-	-	-	-
P ₂	20.80-28.07	14.88	24.16 ± 1.93	-	-	-	-
100-seed weight per plant (g)							
PS (EF)	13.80-17.65	12.24	15.56 ± 0.48	8.86	6.92	61.09	11.15
PS (HY)	12.43-16.30	13.47	14.59 ± 0.22	7.76	7.27	87.80	14.03
SSD	16.47-20.93	11.93	19.40 ± 0.15	6.13	5.97	94.98	11.99
RBP	13.78-16.10	7.76	15.02 ± 0.30	5.04	3.57	50.24	5.21
F ₂	15.93-18.33	7.01	17.64 ± 1.36	-	-	-	-
P ₁	13.57-16.93	11.02	15.20 ± 0.44	-	-	-	-
P ₂	14.60-16.27	5.41	15.25 ± 1.39	-	-	-	-
Seed yield per plant (g)							
PS(EF)	9.00-12.30	15.49	10.28 ± 0.40	9.73	6.80	48.92	9.80
PS(HY)	9.23-11.40	10.52	10.02 ± 0.28	6.54	4.20	41.20	5.55
SSD	9.13-13.87	20.61	11.00 ± 0.77	16.62	11.04	44.16	15.12
RBP	8.93-12.87	18.07	10.28 ± 0.53	11.00	6.20	31.79	7.21
F ₂	5.03-16.73	53.77	10.27 ± 0.98	-	-	-	-
P ₁	8.33-11.67	16.70	9.67 ± 0.75	-	-	-	-
P ₂	8.67-12.17	16.79	9.30 ± 0.79	-	-	-	-
Harvest index (%)							
PS (EF)	32.43-50.84	22.11	42.41 ± 2.97	14.66	7.75	27.99	8.45
PS (HY)	30.03-42.94	17.69	37.47 ± 2.40	13.30	6.88	26.79	7.34
SSD	40.97-67.26	24.29	54.00 ± 4.16	17.50	10.91	38.84	14.00
RBP	34.56-43.85	11.85	38.75 ± 2.75	14.59	7.33	25.23	7.58
F ₂	27.86-86.23	51.16	40.35 ± 6.58	-	-	-	-
P ₁	33.99-48.40	17.49	40.39 ± 3.45	-	-	-	-
P ₂	30.41-46.67	21.09	38.58 ± 1.73	-	-	-	-