



## Research Article

# Combining ability studies for forage quality traits and forage yield in diallel crosses of oats (*Avena sativa* L.)

Mushtaq Ahmad\*, Gul Zaffar, Z.A. Dar, S.M. Razvi, Noorul Saleem and Mehfuza Habib

Division of Genetics and Pant Breeding, S.K. University of Agricultural Sciences and Technology of Kashmir Shalimar campus, Srinagar -191 121 (J&K) India

\* Email: sahilmushtaqdar@rediffmail.com

(Received: 24 Jan 2014; Accepted: 06 Mar 2014)

### Abstract

Studies in oats (*Avena sativa* L.) were carried out to generate information on combining ability (general and specific), nature and magnitude of gene effects and heterosis for forage quality and yield. The experimental material comprised of 10 diverse parents and 45 F<sub>1</sub> crosses (excluding reciprocals) generated through a 10 x 10 diallel mating design. Significant interaction between combining ability effects were observed for all the quality traits and yield. The estimates of additive variance were higher in magnitude than their corresponding dominance variance estimates indicating the role of additive gene action, except green forage yield and dry matter yield. Estimates of GCA effects for crude protein content% indicated that the parent SKO-213 for neutral detergent fibre (NDF), parent SKO-210 for acid detergent fibre (ADF), parent SKO-204 for crude fibre (CF), parent SKO-208 for ash content, parent SKO-207 for green forage yield, parent SKO-212 for dry matter yield were having highly significant and desirable GCA effects for these traits. Cross combinations viz., SKO-212 X SKO-213, SKO-204 X SKO-211, SKO-205 X Sabzaar, Sabzaar X SKO-212 and SKO-205 X SKO-211 exhibited high and desirable sca and heterobeltiosis effects simultaneously for all the quality traits, green forage yield and dry matter yield ha<sup>-1</sup>.

**Key words:** *Avena sativa* L., forage quality and yield, combining ability, heterobeltiosis

### Introduction

Oats (*Avena sativa* L.) is one of the most important winter fodder and grain crops grown throughout world both for animal and human consumption. It is a quick growing, palatable, succulent and nutritious fodder crop which forms an excellent combination when fed along with other cool season legumes. Inadequate supply of quality feed and fodder is the primary cause of lower productivity of milch animals in India (Patel *et al.*, 2011). In Jammu and Kashmir fodder requirement is about 4.31 against the availability of 3.26 million tonnes, there by having a deficient of 1.05 million tonnes on dry matter basis (Anonymous, 2008). Kashmir valley is experienced a long lean period of winter, resulting in scarcity of green and quality fodder which results in drastic decrease in milk and milk production. Therefore, to meet the need of animal products and to maintain good health and potential of livestock in terms of milk, meat and wool, there is a great importance of fodder cultivation to compensate for the fodder scarcity during lean period. The best measure related to forage quality is animal productivity, which can be affected by nutrient intake, digestibility and utilization efficiency. Quality forage must have high intake, digestibility and efficient utilization. Oat genotypes that are low in NDF and ADF have good forage quality because low NDF is associated with high forage intake and

low ADF is associated with high digestibility. Protein content is an important feed factor *per se* with high quality feed having high protein content. The total mineral content of forage is called ash and it represents 3 to 12% DM. The minerals typically determined are calcium and phosphorus.

Combining ability plays a vital role in identifying the potential of lines for obtaining promising segregants. It should pave the way for bringing about a kind of plant type, which could enhance quality and productivity without sacrificing the consumer needs. Productivity and Quality of forage oats can be improved through heterosis breeding. The estimation of heterosis for forage yield and its component characters would be useful to judge the best hybrid combination for exploitation of superior hybrids. However, the selection of promising parents to obtain superior hybrids primarily depends on the predominance of the genes for the additive effect due to heterosis and heterobeltiosis (Beche *et al.*, 2013). Hence the present investigation was carried out with the objectives to study the combining ability, nature of gene action and magnitude of heterosis over better parent for quality and yield traits.

### Material and Methods

The basic material for the present study consisted of 10 diverse genotypes of Oats (*Avena sativa* L.)



viz: SKO-204, SKO-205, SABZAAR, SKO-207, SKO-208, SKO-209, SKO-210, SKO-211, SKO-212 and SKO-213 selected from the germplasm collection maintained at Division of Plant Breeding and Genetics, SKUAST-K, Shalimar. Forty five  $F_1$  crosses (excluding reciprocals) generated through a diallel mating design were evaluated for forage quality and forage yield in a single environment at SKUAST-Kashmir, Shalimar, Srinagar, J&K (India) during *rabi* 2010-2011. The experiment was laid out in a completely randomized block design with three replications. The experimental plot comprised of two rows each of 1m length. Plant spacing was maintained at 30 cm x 10 cm. The forage quality was determined after the samples were dried and crushed to a fine powder. The forage quality parameters for which these genotypes were studied included, Crude protein content (Jackson, 1973), Neutral detergent fibre (Goering and Vansoest, 1970), Acid detergent fibre (Goering and Vansoest, 1970), Ash content % (AOAC, 1984) and crude fibre (Maynard, 1970). Data was subjected to analysis of variance to find significant differences among genotypes for the recorded data. After obtaining the significant differences, data recorded on parents and their  $F_1$  s were subjected to combining ability analysis following Griffing (1956) Method-II and Model-I. The percent increase or decrease of  $F_1$  hybrids over better parent was calculated to estimate the possible heterotic effects for above mentioned parameters (Fonseca and Patterson, 1968).

### Results and Discussion

Significant differences were found among the parents and their crosses for all the traits indicating that the materials selected were diverse and also resulted in creation of substantial genetic variability in the crosses. The contrast of parent's vs hybrids was sizable and highly significant for all traits, pointing to the potential of heterotic effects in improving forage quality and yield traits. The mean sum of squares for general combining ability and specific combining ability were significant for all the traits. It indicated that additive and non additive genetic variance played a significant role in the expression of traits.

The estimates of variance due to additive genetic variance ( $\sigma^2A$ ) were much higher than the corresponding dominance deviation ( $\sigma^2D$ ) for all the quality traits, indicating preponderance of additive gene action as compared to dominance gene action except green forage yield and dry matter yield  $ha^{-1}$  that exhibited non additive gene action. Therefore, preponderance of additive gene action suggested that, methods based on direct selection in early generation such as pedigree method can be applied. Green forage and dry matter yield revealed preponderance of over dominance indicating that the present set of

materials was diverse and contained contrasting alleles which on combination through hybridization increased heterozygosity and could lead to hybrid vigour. These results are general agreement with earlier findings (Prajapati *et al.*, 2010; Akram *et al.*, 2011; Ahmad *et al.*, 2013a) for forage quality and green forage yield.

The estimates of GCA effects listed in Table 2, showed difference of one individual parent to another and from trait to trait. The GCA effects represent the additive nature of gene action. A high general combiner is characterized by its better breeding value when crossed with a number of other parents. Knowledge of the relative importance of additive and non-additive gene action is essential to a plant breeder for the development of an efficient hybridization program. Combining ability analysis helps in the evaluation of pure lines in terms of their genetic value and in the selection of suitable parents for hybridization. Further the GCA variance ( $\sigma^2g$ ) of parents and SCA variance ( $\sigma^2s$ ) of the crosses plays a significant role in the choice of parents.

In the present investigation, the general combining ability effects of 10 lines were estimated to know their genetic worth for use in production of superior progenies. The estimates of GCA effects of parents (Table 2), indicated that the parent SKO-213 (0.468\*\*) and SKO-209 (0.402\*\*) were having highly significant positive GCA effects for crude protein content. For neutral detergent fibre (NDF) parent SKO-210 (-1.886\*\*) and SABZAAR (-1.807\*\*) were having highly significant negative GCA effects. For acid detergent fibre (ADF) parents SKO-204 (-1.262\*\*) and SKO-211 (-0.871\*\*) were having highly significant negative GCA effects. For crude fibre (CF %) parent SKO-208 (-1.175\*\*) and SKO-210 (-1.111\*\*) were having highly significant negative GCA effects. For Ash Content (%) parents SKO-207 (0.433\*\*) and SKO-208 (0.424\*\*) were having highly significant positive GCA effects. For green forage yield parent SKO-212 (6.883\*\*) and SKO-205 (3.654\*\*) were having highly significant positive GCA effects. For dry matter yield parent SKO-212 (4.221\*\*) and SKO-207 (2.389\*\*) were having highly significant positive GCA effects. Parent SKO-207 and SKO-212 with desirable GCA for all traits could be utilised extensively in hybridisation program to accelerate the pace of genetic improvement of quality and forage yield. These genotypes could then serve as a source for isolation of new desirable lines. Parents presenting higher GCA must be preferred to be part of crossing programme, for the selection of promising homozygous lines (Ahmad *et al.*, 2013b). The GCA effects are attributable to additive and additive x additive gene effects. The above mentioned parents have good potential for improving the respective quality traits and may be



used in a multiple crossing program to synthesize an improved population with most of the favourable genes for amelioration of quality and forage yield.

The estimates of specific combining ability effects of the 45 crosses for quality traits, given in Table 3, revealed that out of 45 cross combinations 26 showed desirable significant SCA effect for crude protein content (%). For neutral detergent fibre (%) 21 cross combination exhibited desirable negative SCA effect. For acid detergent fibre out of 45 cross combinations only 23 crosses exhibited desirable significant negative SCA effect. Eighteen and 15 cross combinations exhibited desirable and significant SCA effect for crude fibre and ash content (%). For dry matter yield out of 45 cross combinations only 23 crosses exhibited desirable significant positive SCA effect. For green fodder yield 24 cross combinations possessed desirable and significant SCA effect. Five cross combinations possessed desirable and significant SCA viz; SKO-204 X SKO-211, SKO-205 X SABZAAR, SKO-205 X SKO-211, SABZAAR X SKO-212 and SKO-212 X SKO-213 simultaneously for forage quality, green forage and dry matter yield. The crosses having high specific combining ability (Table 3), for forage quality and yield with other agronomic features need to be evaluated vigorously and selection practiced for isolating desirable transgressive segregants. Parents with positive significant *gca* effects involved additive type of gene action that would be easily fixable. These desirable cross combinations which involve high x low (Table 4) GCA effects in this situation could be utilized in recombination breeding. It is noteworthy that crosses which exhibited consistently positive *sca* effects, it is therefore suggested that *sca* performance may be considered as a criterion for selecting the best crosses. It may also be worthwhile to attempt biparental mating among selected crosses in the advanced generation to permit superior recombination. Parents which have been identified with desirable forage quality parameters could be used in hybridization programmes to improve the quality characteristics in adopted cultivars and desirable cross combinations need to be evaluated vigorously in advanced generation. Crosses which had one good general combiner in their parentage indicated the involvement of additive x dominance type of interaction and they may be advanced for deriving desirable transgressive segregants and homozygous lines in subsequent generations (Ahmad *et al.*, 2013a).

#### Heterosis expression

The estimation of heterosis for quality and forage yield would be useful to judge the best hybrid combination for exploitation of superior hybrids. In the present investigation, the magnitude of heterobeltiosis effects (%) for forage quality, green

forage yield and dry matter yield varied (Table 5). For crude protein content, crosses SKO-212 X SKO-213, SKO-210 X SKO-213 and SKO-209 X SKO-210 exhibited highly significant positive heterobeltiosis values. For NDF, hybrids SKO-205 X SKO-207, SKO-204 X SKO-212 and SKO-205 X SABZAAR showed the desirable values over better parent. For ADF, the crosses SKO-204 X SKO-213, SKO-205 X SKO-207 and SABZAAR X SKO-207 showed significant heterosis over better parent. For crude fibre, the crosses SKO-207 X SKO-210, SKO-212 X SKO-213 and SKO-210 X SKO-211 exhibited highly significant and negative heterobeltiosis values. For ash percentage, heterobeltiosis was positive for most of the crosses. The crosses SKO-205 X SKO-211, SKO-205 X SABZAAR and SKO-212 X SKO-213 showed highly significant and positive heterosis over better parent. For green fodder and dry matter yield  $\text{ha}^{-1}$ , the respective ranges for over better parent heterosis were: -17.906% to 23.908% and -15.667% to 21.333%. The crosses SKO-212 X SKO-213, SKO-204 X SKO-211 and SKO-204 X SKO-208 exhibited significantly positive heterobeltiosis for green fodder and dry matter yield  $\text{ha}^{-1}(\text{q})$ . It is interesting to note that the majority of crosses showing heterobeltiosis and SCA effects for some traits were also among the best performing for the same traits; hence, utilizing heterosis in improving such traits might be rewarding. Furthermore, although quality traits and forage yield are usually reported to be adversely associated (Mohammed and Talib, 2008) some crosses in this study depicted a non adverse association between yield and some quality traits pointing to the possibility of developing crosses with better yield and quality or these cross combinations need further evaluation in segregating generations to identify desirable transgressive segregants. The results from this study indicate that, by using the correct breeding parents and selection procedures, forage quality and yield can be improved.

Variances due to GCA were higher in magnitude than SCA for the forage quality traits which could be exploited for the improvement of these traits by selection. Preponderance of additive type of gene effects suggested directional selection for isolating better homozygous lines from segregating population for these traits. Exploiting heterosis in forage oats to improve quality traits might be promising. Hybrids low in NDF, ADF and CF percentages appears to be attainable without sacrificing high yield levels.

#### **References**

- Ahmad, M., Zaffar, G., Razvi, S.M., Mir, S. D., Rather, M.A. and Dar, Z. A. 2013b. Gene action and combining ability for fodder yield and its attributing traits in oats (*Avena sativa* L.).



- Scientific Research and Essays*, 8(48):2306-2311.
- Ahmad, M., Zaffar, G., Razvi, S.M., Dar, Z.A., Khan, M.H. and Ganie, S.A. 2013a. Combining ability study in oat (*Avena sativa* L.) for physiological, quality traits, forage and grain yield. *African J. Agric. Res.*, 8(43): 5245-5250.
- Akram, Z., Ajmal, S., Khan, K.S., Qureshi, N.R. and Zubir, M. 2011. Combining ability estimates of some yield and quality traits in spring wheat (*Triticum aestivum* L.). *Pakistan J. Bot.*, 43(1): 221-231.
- Anonymous, 2008. Status paper on fodder production in the state of Jammu and Kashmir. Paper presentation 41<sup>st</sup> RCM (Rabi 2008), SKUAST-K, Shalimar, Srinagar, J&K, India.
- AOAC, 1984. Official methods of analysis, 14<sup>th</sup> edn. Arlington, Virginia, USA: Association of Official Analytical Chemists.
- Beche, E., Silva, C.L.D., Pagliosa, E.S., Capelin, M.A., Franke, J., Matei, G., Benin, G. 2013. Hybrid performance and heterosis in early segregants populations of Brazilian spring wheat. *Australian J. Crop Sci.*, 7(1):51-57.
- Fonseca, S. and Patterson, F.L. 1968. Hybrid vigour in a seven parent diallel crosses in common winter wheat (*Triticum aestivum* L.). *Crop Sci.*, 8:85-88.
- Goering, H.K. and Van Soest, P.J. 1970. *Forage fiber analysis* (Apparatus, Reagent. Procedure and some application) Hand book No. 379.ARS, USDA, Washington, D.C., U.S.A.
- Griffing, B. 1956. Concept of general and specific combining ability in relation to dialled crossing systems. *Australian J. Biol. Sci.*, 9: 463 -493.
- Jackson, M.L. 1973. *Soil Chemical Analysis*. Prentice Hall, New Delhi, India.
- Maynard, A.J. 1970. *Methods in Food Analysis*. Academic Press, New York, p.176.
- Mohammed, M.I. and Talib, N.H. 2008. Heterosis and Combining Ability for Quality Traits in Forage Sorghum. *Australian J. Basic and Applied Sci.*, 2(1): 99-104
- Patel, T.U., Arvadia, M.K., Malik, P.K., Patel, D.D. and Patel, P.S. 2011. Production of oats (*Avena Sativa*.L) under different cutting management and split application of nitrogen. *Indian J. Agron.*, 56: 164-167.
- Prajapati, A.S., Vishwakarma, S.R., Vishwakarma, D.N. and Singh, H.P. 2009. Combining ability analysis in intervarietal crosses of oat (*Avena sativa* L.). *Range Manage. Agrofor*, 30(2):163-166.



**Table 1. Analysis of variance for combining ability and estimates of components of variance for forage quality, green forage yield and dry matter yield in oats (*Avena sativa* L.)**

Source of variation	d.f	Mean Square						
		Crude protein content %	Neutral detergent fibre%	Acid detergent fibre %	Crude Fibre%	Ash Content%	Green fodder yield ha <sup>-1</sup> (q)	Dry matter yield ha <sup>-1</sup> (q)
Replication	2	0.0141	0.008	0.238	0.026	0.216	0.209	0.011
Treatment	54	2.825**	47.528**	16.491**	23.601**	110.679**	603.691**	54.345**
Parents	9	7.258**	111.405**	12.078**	9.380**	1.440**	51.134**	79.342**
Hybrids	44	1.734**	35.072**	16.253**	25.420**	19.189**	320.706**	35.987**
Parents Vs Hybrids	1	10.940**	20.670**	66.675**	71.555**	25.056**	8028.050**	9672.889**
Error	108	0.007	9.836	0.107	0.123	0.324	0.313	0.440
GCA	9	108.792*	743.285**	11.304**	81.091**	18.979**	164.318**	324.567**
SCA	45	32.074**	704.223**	4.335**	43.740**	21.913**	208.613**	444.331**
σ <sup>2</sup> A	-	0.710	12.370	4.299	7.597	0.675	27.987	43.23
		±0.011	±1.223	±0.789	±1.314	±0.016	±2.678	±3.779
σ <sup>2</sup> D	-	0.347	7.225	1.878	1.494	0.486	116.876	98.211
		±0.001	±1.113	±0.029	±0.311	±0.032	±3.067	±4.225

\*, \*\* significant at 5 and 1 per cent level, respectively.

**Table 2. General combining ability effects for forage quality, green forage yield and dry matter yield in oats (*Avena sativa* L.)**

Parents	Crude protein content %	Neutral detergent fibre%	Acid detergent fibre %	Crude Fibre%	Ash Content%	Green fodder yield ha <sup>-1</sup> (q)	Dry matter yield ha <sup>-1</sup> (q)
SKO-204	-0.046**	-0.814	-1.262**	0.781**	0.045	1.577**	2.332**
SKO-205	-0.648**	-0.920	-0.821**	0.697**	0.284**	3.654**	1.356**
SABZAAR	-0.787**	-1.807**	0.713**	-0.106	-0.249**	-5.412**	-1.675**
SKO-207	0.100**	-1.669	-0.112*	-0.431**	0.433**	0.573**	2.389**
SKO-208	0.244**	-1.086**	1.641**	-1.175**	0.424**	-0.731**	-3.887**
SKO-209	0.402**	0.544	0.154**	1.408**	0.070	-3.267**	-1.554**
SKO-210	-0.078**	-1.886**	1.229**	-1.111**	-0.108*	-4.188**	-2.332**
SKO-211	0.110**	0.887	-0.871**	-0.492**	-0.316**	1.446**	0.776**
SKO-212	0.235**	-1.044*	-0.826**	-0.303**	0.364**	6.883**	4.221**
SKO-213	0.468**	-1.278*	0.154**	0.731**	0.067	-0.485**	-2.567**
S.E. (g i)	±0.013	±0.095	±0.051	±0.055	±0.050	±0.088	±0.024
S. E. ( gi – gj )	0.020	0.104	0.077	0.082	0.074	0.131	0.213
No of parents showing desirable gea effects	6	5	5	5	4	5	5

\*, \*\* Significant at 5 and 1 per cent level, respectively



**Table 3. Specific combining ability effects for forage quality, green forage yield and dry matter yield in oats (*Avena sativa* L.)**

Crosses	Crude protein content %	Neutral detergent fibre%	Acid detergent fibre %	Crude Fibre%	Ash Content%	Green fodder yield ha <sup>-1</sup> (q)	Dry matter yield ha <sup>-1</sup> (q)
SKO-204 X SKO-205	-1.124**	0.401*	-0.849**	2.186**	-0.292	0.632**	0.363**
SKO-204 X SABZAR	-1.425**	0.278	-2.782**	2.389**	0.141	0.143**	0.243**
SKO-204 X SKO-207	0.488**	-0.988**	-0.057	2.114**	-0.375**	2.285**	1.267**
SKO-204 X SKO-208	0.494**	-1.019**	-0.010	-3.342**	0.461**	4.765**	3.457**
SKO-204 X SKO-209	0.385**	-3.363**	3.676*	3.375**	-0.822**	4.213**	-0.577**
SKO-204 X SKO-210	-0.824**	1.595**	3.701*	-3.906**	-2.400**	-3.899**	-4.897**
SKO-204 X SKO-211	0.387**	-4.503**	-2.799**	-3.525**	4.208**	4.432**	3.556**
SKO-204 X SKO-212	-0.543**	-4.437**	-1.243**	2.453**	4.561**	-1.123**	-2.976**
SKO-204 X SKO-213	0.037	-1.041**	-5.724**	-1.547**	0.225	-0.334**	-0.556**
SKO-205 X SABZAR	1.563**	-4.193**	-4.776*	-3.128**	5.198**	1.568**	0.987**
SKO-205 X SKO-207	-0.610**	-5.383**	-5.299*	1.497**	0.086	1.044**	2.056**
SKO-205 X SKO-208	0.506**	0.053	0.548*	-1.659**	-1.623**	2.324**	3.982**
SKO-205 X SKO-209	0.788**	2.342**	-1.265**	2.958**	0.283	-0.432**	-0.221**
SKO-205 X SKO-210	1.379**	0.901**	1.260**	0.178	4.061**	1.543**	2.445**
SKO-205 X SKO-211	2.340**	-2.391**	-2.540**	-4.742**	5.031**	3.345**	2.045**
SKO-205 X SKO-212	0.955**	0.642**	0.885**	1.931**	1.489**	-0.554**	-1.007**
SKO-205 X SKO-213	0.832**	-2.372**	0.435*	1.036**	-0.114	0.224**	1.788**
SABZAR X SKO-207	0.969**	2.404**	-4.632**	1.733**	-1.181**	2.987**	3.112**
SABZAR X SKO-208	0.125**	0.973**	-0.985**	-0.856**	-1.089**	-3.445**	-2.455**
SABZAR X SKO-209	0.926**	3.229**	-3.199**	-3.649**	-0.583**	1.224**	2.114**
SABZAR X SKO-210	1.317**	4.487**	0.526*	-1.720**	0.694**	1.376**	1.665**
SABZAR X SKO-211	1.135**	-0.305	0.626**	-3.639**	0.602**	-1.234**	-0.556**
SABZAR X SKO-212	1.004**	-3.329**	-2.582**	-0.472**	4.522**	1.879**	2.768**
SABZAR X SKO-213	0.161**	-3.851**	3.401**	1.239**	0.319*	0.980**	1.098**
SKO-207 X SKO-208	-0.942**	-3.498**	1.740**	-0.331	-0.206	3.443**	3.116**
SKO-207 X SKO-209	0.039	0.753**	-0.074	1.186**	0.100	-1.342**	-0.987**
SKO-207 X SKO-210	0.780**	-1.588**	-1.049**	-6.495**	0.777**	1.234**	1.449**
SKO-207 X SKO-211	0.331**	1.120**	-1.102**	-0.114	0.386*	2.231**	2.443**
SKO-207 X SKO-212	-0.753**	1.953**	-3.413**	-7.803**	-0.294	-0.321**	-0.435**
SKO-207 X SKO-213	-0.466**	3.376**	-0.027	1.364**	-0.098	-2.345**	-2.776**
SKO-208 X SKO-209	-0.365**	0.923**	0.773**	-1.670**	-0.108	3.786**	2.980**
SKO-208 X SKO-210	-0.414**	-0.519*	-0.498*	-0.050	3.269**	0.986**	0.994**
SKO-208 X SKO-211	0.187**	-2.011**	-1.102**	1.370**	0.377*	-0.124**	-0.866**
SKO-208 X SKO-212	0.213**	-2.277**	-3.413**	0.941**	-0.697**	-0.008	-0.303**
SKO-208 X SKO-213	-0.430**	0.845**	-0.627**	2.108**	0.094	-1.237**	-2.342**
SKO-209 X SKO-210	-0.572**	-1.263**	-2.115**	-2.634**	-0.425*	0.447**	-1.005**
SKO-209 X SKO-211	0.199**	-2.655**	1.385**	3.278**	0.517**	-0.990**	-1.021**
SKO-209 X SKO-212	-0.796**	1.978**	0.040	2.011**	0.133	-0.248**	-1.225**
SKO-209 X SKO-213	-0.079	0.501*	-0.040	2.658**	-0.100	-0.345**	-1.768**
SKO-210 X SKO-211	-0.280**	0.597**	-1.290**	-5.825**	-2.039**	-4.020**	-3.986**
SKO-210 X SKO-212	0.125*	5.163**	-1.035**	-0.164	-0.719**	-0.704**	-0.342**
SKO-210 X SKO-213	3.412**	0.740**	-2.915**	-3.197**	1.077**	-0.258**	-1.552**
SKO-211 X SKO-212	-0.924**	0.645**	-1.035**	0.369	-0.211	-0.495**	-1.993**
SKO-211 X SKO-213	-0.037	0.367	-0.015	1.425**	-0.114	-0.331**	-1.886**
SKO-212 X SKO-213	4.099**	-3.101**	-0.860**	-3.303**	4.298**	6.469**	4.776**
S. E ( sij)	±0.046	±0.197	±0.174	±0.186	±0.151	±0.024	±0.012
No of crosses showing desirable SCA effects	26	21	23	18	15	24	23



**Table 4. Best parents and crosses identified on the basis of gca and sca effects for forage quality, green forage yield and dry matter yield in oats (*Avena sativa* L.)**

Trait	Parents	GCA	Crosses	SCA	Gca effect of parents
Crude protein content %	SKO-213	0.468**	SKO-212 X SKO-213	4.099**	High x High
	SKO-209	0.402**	SKO-210 X SKO-213	3.412**	Average x High
	SKO-208	0.244**	SKO-205 X SKO-211	2.340**	Average x High
Neutral detergent fibre%	SKO-210	-	SKO-205 X SKO-207	-	Low x Low
	SABZAR	1.886**	SKO-204 X SKO-211	5.583**	Low x Low
	SKO-208	-	SKO-204 X SKO-212	-4.503**	Low x High
		1.807**		-4.437**	
Acid detergent fibre %		-			
		1.086**			
	SKO-204	-	SKO-204 X SKO-213	-5.724**	High x Average
	SKO-211	1.262**	SKO-205 X SABZAR	-4.776**	High x Average
	SKO-212	-	SABZAR X SKO-207	-4.632**	Average x High
Crude Fibre%		0.871**			
		-			
		0.826**			
	SKO-208	-	SKO-207 X SKO-210	-6.495**	High x High
	SKO-210	1.175**	SKO-205 X SKO-211	-4.742**	Average x High
Ash Content%	SKO-211	-	SKO-204 X SKO-210	-3.696**	Average x High
		1.111**			
		-			
		0.492**			
	SKO-207	0.433**	SKO-205 X SABZAR	5.198**	High x Average
Green fodder yield ha <sup>-1</sup> (q)	SKO-208	0.424**	SKO-211 X	5.031**	Average x Average
	SKO-212	0.364**	SABZAR	4.561**	Low x High
			SKO-204 X SKO-212		
	SKO-212	6.883**	SKO-212 X SKO-213	6.469**	High x Average
	SKO-205	3.654**	SKO-204 X SKO-208	4.765**	High x Average
Dry matter yield ha <sup>-1</sup> (q)	SKO-204	1.557**	SKO-204 X SKO-211	4.432**	High x High
	SKO-212	4.221**	SKO-212 X SKO-213	4.776**	High x Average
	SKO-207	2.389**	SKO-205 X SKO-208	3.982**	High x Average
	SKO-204	2.332**	SKO-204 X SKO-211	3.566**	High x High



**Table 5. Magnitude of heterosis over better parent % (heterobeltiosis) for forage quality, green forage yield and dry matter yield in oats (*Avena sativa* L.)**

Crosses	Crude protein content %	Neutral detergent fibre%	Acid detergent fibre %	Crude Fibre%	Ash Content%	Green fodder yield ha <sup>-1</sup> (q)	Dry matter yield ha <sup>-1</sup> (q)
SKO-204 X SKO-205	-2.334**	2.321*	-2.765**	5.332**	-3.004**	5.098**	5.446**
SKO-204 X SABZAR	-5.556**	2.578**	-4.443**	5.567**	0.007	6.880**	5.334**
SKO-204 X SKO-207	3.567**	-2.989**	-0.341	6.443**	-3.889**	11.233**	9.007**
SKO-204 X SKO-208	3.998**	-3.008**	-0.112	-7.224**	4.807**	17.899**	15.678**
SKO-204 X SKO-209	2.978**	-3.045**	5.775**	8.552**	-03.090**	16.775**	-9.002**
SKO-204 X SKO-210	-3.775**	3.678**	5.332**	-7.665**	-6.001**	-14.643**	-13.089**
SKO-204 X SKO-211	3.008**	-6.556**	-4.577**	-8.212**	9.078**	18.467**	14.342**
SKO-204 X SKO-212	-4.908**	-6.867**	-3.221**	5.345**	8.988**	-8.046**	-12.333**
SKO-204 X SKO-213	1.876**	-3.443**	-7.553**	-4.589**	0.045	-8.870**	-6.774**
SKO-205 X SABZAR	6.776**	-6.774**	-6.332*	-7.890**	13.034**	9.003**	6.773**
SKO-205 X SKO-207	-3.089**	-7.443**	-7.367*	4.579**	0.001	10.886**	11.233**
SKO-205 X SKO-208	4.076**	1.236**	2.362**	-4.887**	-5.445**	12.446**	12.332*
SKO-205 X SKO-209	5.664**	4.455**	-3.876**	5.656**	0.020	-6.776**	-7.001**
SKO-205 X SKO-210	5.334**	2.116**	3.559**	0.131	9.040**	9.880**	10.989**
SKO-205 X SKO-211	7.453**	-4.393**	-4.576**	-8.778**	13.980**	14.080**	13.132**
SKO-205 X SKO-212	1.221**	2.665**	2.788**	4.223**	4.556**	-5.556**	-4.908**
SKO-205 X SKO-213	2.643**	-4.255**	2.576**	4.558**	-0.030	6.673**	4.553**
SABZAR X SKO-207	3.997**	4.587**	-6.787**	4.009**	-4.087**	11.212**	10.989**
SABZAR X SKO-208	5.977**	2.006**	-2.344**	-3.908**	-5.067**	-14.334**	-13.432**
SABZAR X SKO-209	3.334**	5.789**	-5.221**	-7.880**	-3.332**	8.009**	9.998**
SABZAR X SKO-210	6.667**	6.543**	2.322*	-4.776**	4.221**	9.012**	8.908**
SABZAR X SKO-211	7.445**	-2.334**	2.978**	-7.856**	3.007**	-10.880**	-9.008**
SABZAR X SKO-212	3.443**	-5.312**	-4.445**	-3.070**	9.889**	9.030**	7.887**
SABZAR X SKO-213	1.997**	-5.745**	5.887**	4.765**	4.005*	5.001**	3.332**
SKO-207 X SKO-208	-3.233**	-5.887**	3.443**	-0.004	-0.304	14.231**	12.890**
SKO-207 X SKO-209	1.001**	2.643**	-0.221	8.347**	0.004	-15.887**	-14.143**
SKO-207 X SKO-210	2.221**	-3.221**	-1.049**	-14.562**	3.114**	8.900**	7.664**
SKO-207 X SKO-211	1.556**	3.665**	-3.775**	-0.221	4.223*	11.070**	9.003**
SKO-207 X SKO-212	-5.665**	3.451**	-5.769**	-10.800**	-0.090	-5.676**	-4.444**
SKO-207 X SKO-213	-3.883**	5.424**	-1.002	4.021**	-0.098	-11.898**	-10.122**
SKO-208 X SKO-209	-0.365**	2.774**	2.331**	-4.443**	-3.009**	4.445**	5.344**
SKO-208 X SKO-210	-7.665**	-2.089**	-2.557**	-0.009	8.885**	5.989**	4.332**
SKO-208 X SKO-211	2.113**	-4.464**	-3.115**	4.080**	3.060*	-6.778**	-5.554**
SKO-208 X SKO-212	1.231**	-4.806**	-5.881**	3.907**	-4.505**	-0.010	-0.033
SKO-208 X SKO-213	-3.332**	2.327**	-2.115**	5.076**	0.102	8.777**	-8.889**
SKO-209 X SKO-210	-6.789**	-3.007**	-4.221**	-5.089**	-3.065**	5.006**	-6.443**
SKO-209 X SKO-211	1.087**	-4.440**	3.334**	7.988**	4.557**	-6.770**	-5.443**
SKO-209 X SKO-212	-3.345**	3.009**	0.002	5.770**	0.990*	-5.225**	-3.324**
SKO-209 X SKO-213	-1.876**	2.655**	-0.223*	5.606**	-0.050	-6.664**	-7.876**
SKO-210 X SKO-211	-3.112**	2.703**	-3.229**	-11.990**	-6.774**	-17.906**	-15.667**
SKO-210 X SKO-212	2.434*	8.060**	-3.677**	-3.006**	-4.558**	-5.778**	-2.334**
SKO-210 X SKO-213	9.341**	2.670**	-4.553**	-7.884**	6.080**	-6.993**	-7.080**
SKO-211 X SKO-212	-3.467**	2.909**	-3.332**	0.006	-0.012	-5.552**	-2.098**
SKO-211 X SKO-213	-1.589**	2.507**	-0.115	4.063**	1.033*	-6.996**	-5.554**
SKO-212 X SKO-213	10.569**	-5.967**	-2.235**	-12.900**	9.998**	23.908**	21.333**
S.E.	0.0981	0.0572	0.0411	0.0890	0.0465	0.0778	0.0557
C.D. (5%)	0.5476	0.2671	0.3013	0.2029	0.1041	0.3445	0.3687