


Electronic Journal of Plant Breeding



Research Article

A novel high yielding dual-purpose sorghum variety GDJ 1 (Banas Surya) for semi-arid region of Gujarat

R. A. Gami*, R. N. Patel, P. R. Patel and S. K. Jain

Sorghum Research Station & AICRP Sorghum,
Sardarkrushinagar Dantiwada Agricultural University, Deesa-385 535, Gujarat, India
*E-Mail: ramangami@gmail.com

Abstract

The new sorghum variety GDJ 1 (Banas Surya) breed through pedigree selection method from the cross between SPV 2113 × GFS 5 at Sorghum Research Station, Sardarkrushinagar Dantiwada Agricultural University, Deesa, Gujarat. The variety GDJ 1 was tested in a preliminary evaluation trial (PET) in *kharif* 2016. It was found promising and further it was tested in state multilocation trials from *kharif* 2017 to 2019. It was also tested under the AICRP-sorghum IVT trial in *kharif* 2018 as SPV 2565 at fifteen locations (Zone-I, II and III) across the nation. The sorghum variety GDJ 1 was tested under a total of 20 state trials against various check varieties *viz.*, CSV 20, GJ 39, GJ 42, GDJ 1 and GJ 43. The mean performance of GDJ 1 for grain yield over 21 locations, including the preliminary trial was 2506 kg/ha. In the case of dry fodder yield, the mean performance over 22 location, including the preliminary trial was 167.0 q/ha. In North Gujarat, based on seven testing trials GDJ 1 exhibited high grain yield (2576 kg/ha) with an increment of 50.0, 50.7, 19.6, 21.5 and 41.8 per cent, respectively and dry fodder (228.7 q/ha) with an increment of 49.8, 52.3, 10.9, 49.6 and 34.2 per cent over the years and locations against checks GJ 39, GJ 42, GJ 43, GNJ 1 and CSV 20, respectively. It matures within 100-105 days, tall in nature, very long and broad leaves, the long symmetric semi-loose panicle have a medium neck with long branches and medium glume length. The grain is medium bold and yellow-white. From the quality point of view the released variety have a fair amount of protein in grain (10.1%) and dry fodder (9.20%) and low tannin (0.215 mg/g) in grain indicated the good quality of grain and dry fodder. It is moderately resistant to diseases like leaf blight, anthracnose, grain mold, ergot and lower infestation of shoot fly and stem borer.

Key words

Sorghum, variety, grain yield, dry fodder yield and morphological traits.

INTRODUCTION

Sorghum or Great millet [*Sorghum bicolor* (L.) Moench] is the most important food and fodder crop of the semi-arid tropics. Sorghum grains are essential as food for human being and livestock feed. In Gujarat, only grain type sorghum occupies an area of 73.6 thousand hectares with an annual production of 1.04 lakh tonnes with a productivity of 1419 kg/ha. The fodder area approximately more than 5.0 lakh hectares. Sorghum mostly is grown as dual and fodder purpose in Gujarat. The animal husbandry business is increasing, so the sorghum has an excellent opportunity to fulfil the shortage of feed and fodder. Sorghum is an important crop of semi-arid tropics in Asia

and Africa. India is the 2nd leading producer of sorghum after the USA and the other sorghum producing countries *viz.*, Nigeria, China, Mexico, Sudan and Argentina.

For any agriculture-based country, fodder crops are as much essential as grain crops. The livestock sector being a sub-sector of Indian agriculture contributes about 9 per cent to GDP and employs around 8 per cent of the labour power. In such a context, crops with multiple uses fetch more area as compared to others because crop residues used as animal feed and fodder. Experiential studies in India have revealed that enhancing the quality and

quantity of feed input has a more significant effect than breed improvement on increasing milk productivity (Tonapi *et al.*, 2011). India is anticipating for the white revolution, which is possible only with an adequate supply of nutritious feeds and fodders. The *livestock population in India* is nearly 512.05 million in 2019 and they produced about 165.4 million tonnes of milk annually (Anonymous, 2017-18). There are many cereal fodder crops like sorghum, maize, pearl millet for *kharif* season. High productivity, fast-growing habit and better water use efficiency under drought make it a vital forage crop of semi-arid tropics. It is extremely edible and more digestible than maize and pearl millet and uses approximately 40-50 per cent less water than corn to produce the same dry matter. As far as forage quality is concerned, it is defined as an expression of characteristics that affect consumption and is assessed by digestibility and palatability (Amigot *et al.*, 2005). The demand for dual sorghum is higher in Gujarat. Sorghum is grown in almost all the districts of Gujarat to meet the grain and fodder requirement for maintaining the milch animals. The demand for dual type sorghum increasing day by day in the semi-arid region of Gujarat as it valued both for its grain as well as for its excellent fodder.

In-plant breeding, improvement of a crop concerning yield and specific traits is a continuous programme and the improved varieties in terms of productivity and resistance to biotic and abiotic stress by reformed genetic background are advanced and released for the benefit of the farming community is the prime motto of this centre. At present, cultivars GJ 42 and GNJ 1 for grain, GJ 39, GJ 43 for dual

purpose are cultivating. The local varieties like *malwan* have demerits like a high infestation of pest, late maturity and poor yields both for grain and fodder. Therefore, it is a need to develop a variety having high grain as well as fodder yield potential with early to medium maturity. The released variety GDJ 1 was developed from cross SPV 2113 × GFS-5 followed by continuous evaluation and selection.

MATERIAL AND METHODS

A new sorghum variety GDJ 1 breed through pedigree selection methods from cross SPV 2113 × GFS-5 at Sorghum Research Station, Sardarkrushinagar Dantiwada Agricultural University, Deesa, Gujarat with the name of DS 172. The cross was made in summer 2009 followed by pedigree selection was done from *kharif*, 2009 to 2015. This variety has advanced with main objectives were dual-purpose, *i.e.* high dry fodder yield as well as a fair amount of grain yield. The variety GDJ 1 was tested in a preliminary evaluation trial (PET) in *kharif*, 2016. It was found promising and further, it was tested in state multilocation trial from *kharif* 2017 to 2019. It was also tested under the AICRP sorghum- IVT trial in *kharif* 2018 as SPV 2565 at fifteen locations (Zone-I, II and III) across the nation. The culture was also screened for disease and pest under field condition at state as well as in AICRP trials (Anonymous 2018-19). The DNA fingerprinting of variety GDJ 1 along with four checks (GJ 43, GJ 39, GNJ 1 and CSV 20) were performed by using 11 ISSR primers (Table 1).

Table 1. List of primers used in fingerprinting

Sr. No.	Name of primer	Sequence of primer	Sr. No.	Name of primer	Sequence of primer
1	ISSR 1	CACACACACACACACAGT	7	ISSR 16	CACACACACACACACAAA
2	ISSR 2	CACACACACACACACAGC	8	ISSR 21	GCCTCTCTCTCTCTCT
3	ISSR 3	CACACACACACACACAGG	9	ISSR 22	GACTCTCTCTCTCTCT
4	ISSR 4	CACACACACACACACAGA	10	ISSR 23	GTCTCTCTCTCTCTCT
5	ISSR 7	CACACACACACACACATG	11	ISSR 25	GCCCTCTCTCTCTCTCT
6	SSR 8	CACACACACACACACATA			

RESULTS AND DISCUSSION

The sorghum variety GDJ 1 was evaluated in PET during *kharif* 2016 against various check varieties *viz.*, CSV 20 and GJ 39. The tested genotype exhibited a significantly high grain yield (3554 kg/ha) with an increment of 53.9 and 58.0 per cent over checks CSV 20 and GJ 39, respectively (Table 2). Similarly, in the case of dry fodder yield, the variety GDJ 1 exhibited significantly high dry fodder yield (215.8 q/ha) with an increment of 21.0 and 39.4 per cent over checks CSV 20 and GJ 39, respectively (Table 3).

The sorghum variety GDJ 1 was tested under a total of 20 state trial during *kharif* 2017-2019 against various check

varieties *viz.*, CSV 20, GJ 39, GJ 42, GNJ 1 and GJ 43. The mean performance of GDJ 1 for grain yield over 21 locations, including the preliminary trial was 2506 kg/ha. In the case of dry fodder yield the mean performance over 22 locations, including the preliminary trial was 167.0 q/ha. Based on the comparative mean analysis, it produced 12.4, 46.2 and 9.2 per cent higher grain yield against the check CSV 20, GJ 39 and GJ 42, respectively (Table 2). In the case of dry fodder, it produced 23.0, 36.7, 26.8, 29.0 and 8.2 per cent higher dry fodder yield against the check CSV 20, GJ 39, GJ 42, GNJ 1 and GJ 43, respectively (Table 3). In North Gujarat based on seven testing trials, GDJ 1 exhibited high grain yield (2576 kg/ha) with an

increment of 50.0, 50.7, 19.6, 21.5 and 41.8 per cent, respectively (Table 2) and dry fodder (228.7 q/ha) with an increment of 49.8, 52.3, 10.9, 49.6 and 34.2 per cent over

the years and locations against checks GJ 39, GJ 42, GJ 43, GNJ 1 and CSV 20, respectively (Table 3).

Table 2. Grain yield performance of sorghum variety GDJ 1 in comparison with check varieties in the Gujarat state

Year/ & Season	Trial	Location	Grain yield (kg/ha)					S.Em (±)	CD at 5 %	CV %	
			GDJ 1	CSV 20 (NC) a	GJ 39- DS (LC) b	GJ 42 (LC) c	GNJ 1 (LC) d				GJ 43-DS- (LC) e
Kharif 2016	PET	Deesa	3554 ^{ab}	2310	2250			221.9	650.5	16.0	
		MEAN (1)	3554	2310	2250						
		% increase over the check	53.9	58.0							
2017 Kharif	SSVT	Surat	2364 ^b	2045	1564	2128	2218	190.0	544.0	15.9	
		Dediyapada	3723 ^{ab}	3006	1919	3345	3495	207.0	593.0	12.5	
		Mangrol	2703 ^b	2443	1714	2754	2809	247.0	708.0	17.3	
		Achhaliya	4050 ^{ab}	2272	1450	3687	3364	349.0	999.0	16.8	
		Viramgam	1278 ^a	708	1063	1220	1543	91.0	260.5	13.3	
		MEAN (5)	2824	2095	1542	2627	2686	-	-	-	
% increase over the check	34.8	83.1	7.5	5.1							
Kharif 2018	LSVT	Deesa	3999 ^{abcd}	2664	3071	2979	3309	218.0	629.0	11.4	
		Bhiloda	2826 ^{bc}	2480	1597	1550	2443	159.0	458.0	12.0	
		Viramgam	949	1949	1971	2719	2895	140.0	405.0	9.7	
		Surat	3223 ^{ab}	2369	2275	2816	4090	294.0	849.0	14.7	
		Waghai	4765 ^{abcd}	3704	4090	3885	4106	167.0	482.0	7.1	
		Mangrol	1235	2075	1602	1768	2141	130.0	375.0	12.6	
		Achhaliya	1208 ^b	2863	841	1051	1812	76.0	220.0	8.9	
		Dediyapada	4074 ^{abd}	3362	1923	3420	3216	229.0	661.0	11.4	
		MEAN (8)	2785	2683	2171	2524	3002	-	-	-	
% increase over the check	3.8	28.3	10.4								
Kharif 2019	LSVT	Deesa	1360	847			1188	1271	147.0	427.0	23.8
		Bhiloda	1919 ^{ad}	1376			1551	1626	95.0	277.0	10.4
		Aseda	1800 ^{ade}	1227			1306	1348	102.0	295.0	12.6
		Surat	1782	2528			3190	2255	213.0	618.0	15.1
		Waghai	2382	2600			2279	2621	107.0	312.0	7.5
		Mangrol	2287	2261			2901	2086	229.0	666.0	15.1
		Achhaliya	1150	1743			2546	1592	161.0	467.0	13.4
		MEAN (7)	1811	1797			2137	1828	-	-	-
% increase over the check	0.8										
Comparative mean over the check CSV 20 (21)			2506	2230	-	-	-	-	-	-	
Comparative mean over the check GJ 39 (14)			2854	-	1952						
Comparative mean over the check GJ 42 (13)			2800	-	-	2563					
Comparative mean over the check GNJ 1 (20)			2454	-	-	-	2630				
Comparative mean over the check GJ 43 (7)			1811	-	-	-	-	1828			
Overall % increase over the checks in all over Gujarat state			-	12.4	46.2	9.2	-	-			
Overall % increase over the checks in north Gujarat zone (semi-arid)			2576	41.8	50.0	50.7	21.5	19.6			
Frequency of top non-significant groups			10/21	2/21	0/14	3/13	5/20	2/7			

Where: Figure in parenthesis indicates the number of locations; a, b, c, d and e indicate significantly superior to respective check variety for grain yield

Table 3. Dry fodder yield performance of proposed sorghum variety GDJ 1 in comparison with check varieties in the Gujarat state

Year/ Season	Name of Trial	Location	Dry fodder yield (q/ha)					S. Em (±)	CD at 5 %	CV %	
			GDJ 1	CSV 20 (NC) a	GJ 39- DS (LC) b	GJ 42 (LC) c	GNJ 1 (LC) d				GJ 43- DS- (LC) e
Kharif 2016	PET	Deesa	215.8 ^{ab}	178.3	154.8			12.7	36.1	11.9	
		MEAN (1)	215.8	178.3	154.8						
		% increase over the check	21.0	39.4							
Kharif 2017	SSVT	Surat	131.9	112.3	106.4	126.9	112.3	9.5	NS	13.5	
		Dediyapada	248.2 ^{abcd}	145.0	212.8	190.3	177.3	5.0	14.3	5.4	
		Mangrol	93.0	138.7	91.0	136.7	143.8	8.6	24.6	12.7	
		Achhaliya	104.4	96.5	96.5	89.4	94.6	6.6	18.8	11.7	
		Viramgam	147.0 ^b	119.0	110.0	137.0	147.0	9.9	28.3	14.5	
		MEAN (5)	144.9	122.3	123.4	136.1	135.0	-	-	-	
% increase over the check	18.5	17.5	6.5	7.3							
Kharif 2018	LSVT	Deesa	338.8 ^{abcd}	244.3	207.5	207.5	228.5	18.1	52.3	13.2	
		Bhiloda	219.4 ^{abcd}	153.0	154.5	159.1	152.2	8.4	24.3	9.3	
		Viramgam	286.7 ^{bcd}	244.3	207.7	202.0	161.3	17.7	51.0	15.0	
		Surat	192.3 ^{abcd}	105.3	114.8	112.7	126.6	10.2	29.5	13.7	
		Waghai	117.9 ^{abcd}	89.9	86.9	88.4	91.6	5.6	16.2	10.9	
		Mangrol	54.3	62.5	44.8	52.7	63.2	5.9	17.0	16.8	
		Achhaliya	93.1	106.4	80.0	108.5	122.6	4.5	13.0	7.7	
		Dediyapada	102.2 ^{abcd}	62.8	47.8	68.3	64.9	5.2	14.9	12.2	
		MEAN (8)	175.6	133.6	118.0	124.9	126.4	-	-	-	
		% increase over the check	31.5	48.8	40.6	39.0					
Kharif 2019	LSVT	Deesa	185.2 ^{ad}	115.7			113.1	167.2	11.7	34.1	14.0
		Bhiloda	188.9 ^{ad}	158.1			122.2	166.4	7.9	23.1	9.0
		Aseda	223.8 ^{ade}	172.6			156.9	205.3	13.5	39.0	13.5
		Surat	172.3 ^d	182.6			132.4	171.0	13.2	38.2	13.7
		Waghai	93.4	102.2			93.0	99.7	4.5	13.1	8.3
		Mangrol	87.8	90.2			77.3	94.3	6.6	19.2	13.5
		Achhaliya	202.9 ^d	195.0			165.1	188.0	11.5	33.5	11.2
		Viramgam	175.7 ^{ade}	113.7			135.0	137.7	8.2	23.9	9.7
		MEAN (8)	166.3	141.3			124.4	153.7	-	-	-
		% increase over the check	17.7	33.7	8.2						
Comparative mean over the check CSV 20 (22)			167.0	135.8	-	-	-	-	-	-	
Comparative mean over the check GJ 39 (14)			167.5	-	122.5	-	-	-	-	-	
Comparative mean over the check GJ 42 (13)			163.8	-	-	129.2	-	-	-	-	
Comparative mean over the check GNJ 1 (21)			164.7	-	-	-	127.7	-	-	-	
Comparative mean over the check GJ 43 (8)			166.3	-	-	-	-	153.7	-	-	
Overall % increase over the checks in all over Gujarat state			-	23.0	36.7	26.8	29.0	8.2			
Overall % increase over the checks in north Gujarat zone (semi-arid)			228.7	34.2	49.8	52.3	49.6	10.9			
Frequency of top non-significant groups			16/22	9/22	1/14	2/13	6/21	5/8			

Where: Figure in parenthesis indicates the number of locations;
a, b, c, d and e indicates significantly superior to respective check variety for dry fodder yield

The variety GDJ 1 was screened for yield performance in IVT with the name SPV 2565 in sixteen locations of Zone I, II and III during *kharif* 2018. The mean performance GDJ 1 among fifteen trials of IVT, it showed 25.7, 12.8 and 19.9 per cent grain yield advantage against CSV 17 in

Zone I, II and III, respectively (**Table 4**). For fodder yield, it showed 45.8, 31.1 and 88.6 per cent yield advantage against CSV 17 in Zone I, II and III, respectively (**Table 5**).

Table 4. Grain yield of proposed sorghum variety GDJ 1 in an initial varietal trial in the year *Kharif* 2018 at different Zones of India

Locations	Grain yield (kg/ha)			CD at 5%	CV%	
	GDJ 1	CSV 17 a	CSV 20 b			
Zone I	Chamarajanagar	4148	3667	4407	1247	18.8
	Coimbatore	3774 ^{ab}	1541	2017	1208	20.0
	Dharwad	4171	4609	4981	939	12.1
	Hagari	3234	3590	5472	1482	21.1
	Palem	4946 ^a	2715	4750	1451	21.9
	Nandyal [#]	1894	1557	2971	1115	30.7
	Mean (5)	4055	3224	4325	1010	18.8
	% Increase over the checks	25.7	-			
Zone II	Akola	3093	3008	3855	652	10.9
	Bhulandnagar	2870	2433	3211	977	20.3
	Indore	3592	2496	3492	918	16.6
	Parbhani	2014	1963	2244	515	13.7
	Washim	3116	3122	3781	734	12.3
	Mean (5)	2937	2604	3317	502	14.9
	% Increase over the checks	12.8	-			
Zone III	Deesa	2489 ^a	649	2341	818	21.0
	Surat	3075	2318	2672	854	17.3
	Udaipur	2430	1657	3840	1217	20.9
	Viramgam	1508	3301	2361	668	16.2
	Mean (5)	2376	1981	2804	1027	19.4
	% Increase over the checks	19.9	-			
All India Mean	3176	2648	3530	491.5	18.09	
	% Increase over	19.9	-			

Data not considered due to high CV %,

a and b indicate significantly superior to respective check variety

Yield with quality is also of prime importance as the sorghum is being used grain as well as fodder purpose. So that it satisfies the need of both human for grain and dry fodder for animal consumption in semi-arid regions. The grain colour of released cultivar GDJ 1 is yellow-white which may be more attractive among farmers for fetching higher market price in this region and consumers point of view. The GDJ 1 have a 10.10 per cent crude protein which was higher against check cultivars viz., CSV 20 (7.8 %), GJ 43 (8.1 %) and GNJ 1 (7.5%). It has a low tannin content (0.215 mg/g) as compared to check varieties. The high tannin gives a bitter taste in some foods and also decline protein digestibility and feed efficiency in human and livestock. On the other hand, the released cultivar GDJ 1 has a high 0.165 (%) tryptophan as compared to

check varieties viz., CSV 20 (0.134 %), GJ 43 (0.117%) and GNJ 1 (0.119%). The crude protein content and crude fiber of dry fodder are one of the essential components for fodder quality in sorghum. The sorghum released variety GDJ 1 was found to better concerning the crude protein content (9.2%) and crude fiber (24.24 %), which was higher against check cultivars in order to crude protein and fiber content viz., CSV 20 (8.2 % & 15.66), GJ 43 (7.9 & 22.39 %) and GNJ 1 (8.4 & % 20.31) (**Table 6**).

The released cultivar GDJ 1 having an attractive suture in field view (**Fig. 1**). It has yellow, green seedling anthocyanin, leaf sheath anthocyanin and leaf midrib coloration. It is tall, and it attained an average of 274.4 cm height in normal condition, very long (>80 cm) and

Table 5. Dry fodder yield performance of sorghum variety GDJ 1 in an initial varietal trial in the year *Kharif* 2018 at different Zones of India

Locations	Dry fodder Yield (q/ha)			CD at 5%	CV%	
	GDJ 1	CSV 17 (a)	CSV 20 (b)			
Zone I	Chamarajanagar	86.1	93.3	79.6	29.1	19.1
	Coimbatore	83.9a	49.2	80.5	21.5	17.5
	Dharwad	109.9a	48.5	118.2	23.2	11.0
	Hagari	101.9	99.5	108.8	23.6	14.0
	Nandyal	129.6a	60.2	175.9	43.5	20.2
	Palem#	135.9	85.6	114.1	161.0	79.3
	Mean	102	70	113	24.3	16.7
	% Increase over the checks	45.8	-			
Zone II	Akola	142.7a	120.3	139.4	10.3	4.6
	Bhulandnagar	127.3	112.6	130.6	23.4	10.6
	Indore	129.6a	44.4	129.6	35.6	18.9
	Parbhani	118.7	104.3	100.7	21.9	11.9
	Washim	143a	122.9	138.9	8.6	3.8
	Mean	132	101	128	16.9	10.7
	% Increase over the checks	31.1	3.5			
Zone III	Deesa	318.5ab	122.2	292.6	61.8	14.4
	Surat	65.2	65.9	77.3	25.3	15.9
	Udaipur	157.5	81.7	151.3	50.8	20.5
	Viramgam	167.5	105.9	165.2	31.2	12.2
	Mean	177	94	172	46.9	16.4
	% Increase over the checks	88.6	3.2			
All India Mean	134	88	134	16.67	15.17	
	% Increase over the checks	53.2	0.7			

Data not considered due to high CV %;

a and b indicate significantly superior to respective check variety

Table 6. Biochemical parameters of variety GDJ 1 (Banas Surya) with checks for grain and dry fodder

Sr. No.	Quality parameters	GDJ 1	CSV 20 (NC)	GJ 43 (C)	GNJ 1 (C)
Grain					
1	Moisture (%)	10.29	10.64	11.16	10.71
2	Crude Protein (%)	10.1	7.8	8.1	7.5
3	Fat (%)	3.6	3.5	3.6	3.5
4	Ash (%)	1.85	1.86	1.67	1.71
5	Crude fiber (%)	0.96	1.37	0.90	1.10
6	Starch (%)	50.7	51.5	51.5	57.0
7	Tannin (mg/g)	0.215	0.232	0.244	0.231
8	Phenol (g)	0.154	0.142	0.133	0.127
9	Lysine (%)	0.493	0.532	0.533	0.628
10	Tryptophan (%)	0.165	0.134	0.117	0.119
Dry Fodder					
1	Moisture (%)	12.0	10.7	11.8	11.3
2	Ash (%)	5.60	6.54	5.38	8.31
3	Crude fiber (%)	24.24	15.66	22.39	20.31
4	Crude Protein (%)	9.2	8.2	7.9	8.4



Fig.1: Field view of sorghum variety GDJ 1 (DS 172) BANAS SURYA

broadleaf blade (6.1 - 8.0). The GDJ 1 have a long panicle length (31-40 cm) with long branches (10.1-15 cm). It has semi-loose panicle compactness devoid of lemma arista

formation and circular and attractive medium bold grain, with medium glume length (**Table 7 and Fig. 2 and 3**).

Table 7. Morphological characters of proposal variety GDJ 1 with check GJ 43 (As per DUS guidelines)

S. N.	Characters	GDJ 1 (DS 172)	GJ 43 (C)
1	Seedling: anthocyanin coloration	Yellow-green	Yellow-green
2	Leaf-sheath anthocyanin coloration	Yellow-green	Yellow-green
3	Leaf midrib colour	Yellow-green	Yellow-green
4	Plant: time of panicle emergence (50% of the plants with 50% anthesis)	Medium (66-75 days)	Medium (66-75 days)
5	Flag leaf yellow coloration of the midrib	Present	Present
6	Lemma arista formation	Absent	Absent
7	Plant total height	Long (226 - 300 cm)	Long (Tall)
8	Stem diameter (girth)	Medium (2 - 4 cm)	Medium (2 - 4 cm)
9	Leaf: length of the blade	Very Long (> 80 cm)	Long (61 – 80 cm)
10	Leaf : width of blade	Broad (6.1 - 8.0)	Broad (6.1 - 8.0)
11	Panicle length (without peduncle)	Long (31-40 cm)	Medium (21-30 cm)
12	Ear head compactness	Semi loose	loose
13	Panicle shape	Symmetric	Symmetric
14	The neck of panicle (visible length above sheath)	Medium (10.1 - 15 cm)	Medium (10.1 - 15 cm)
15	Length of branches (middle third of panicle)	Long (10.1-15 cm)	Medium (5.1-10 cm)
16	Glume length (% grain covered)	Medium (75 %)	Short (50 %)
17	Length of flower	Medium	Medium
18	Dry anther colour	Greyed orange	Greyed orange
19	Anther length	Short (<3.0 mm)	Short (<3.0 mm)
20	Stigma : Anthocyanin coloration	Absent	Absent
21	Stigma :Yellow coloration	Present	Absent
22	Stigma length	Medium (1-2 mm)	Short (<1mm)
23	Grain shape (dorsal and profile view)	Circular	Elliptic
24	Caryopsis colour after threshing	Yellow white	Yellow white
25	Size of a mark of germ	Medium	Medium
26	Threshability	Freely threshable	Freely threshable
27	Grain weight (1000 grain wt. in g)	Medium (26 - 35 g)	Medium (26 - 35 g)

During the cropping season, sorghum crop experiences many diseases viz., leaf blight, anthracnose, grain mold and ergot. In the case of disease, grain mould considered as the most important biotic factor that constrains the production as well as the quality of grain sorghum worldwide (Thakur *et al.*, 2006). On the other hands, major pests like shoofly and stem borer, which harshly distress to sorghum crop (Thakur *et al.*, 2019). Shoot fly is one of the disparaging pests and causes bare damage in sorghum at the time of the early seedling stage (7-30 days of seedling). While *C. partullus* (stem borer) is also playing a significant negative role in the decrease of sorghum production in the country. The August month

is preferable in which maximum infestation (about 4-45%) occurs, but it declined gradually in later months (Singh *et al.*, 1985). The released cultivar tested in various locations in AICRP trials as well as an in-state trial for evaluating reaction toward major disease and pest. Under field condition, the cultivar GDJ 1 gave a moderately resistance reaction toward leaf blight, anthracnose, grain mold and ergot diseases (**Table 8 & 9**). It showed lesser infestation of shoot fly (29.00%) and stem borer dead heart (31.27 %) in state trials (**Table 10**). Under the AICRP trial, this cultivar also showed moderate to lesser reaction toward shoot fly (57.2%) and stem borer dead heart (19.3 %) (**Table 11**).




Fig. 2. Important DUS characteristics of variety GDJ 1(DS 172)



Stigma : Anthocyanin colouration : Absent
Yellow colouration : Present
Length : Medium (1-2 mm)



Leaf; length of blade: Very long (>80cm)
Width of blade: Broad (6.1-8.0 cm)



Panicle density : Semi loose
Panicle Shape : Symmetric
Panicle length : Long



Neck of panicle (visible length above sheath) : Medium (10.1-15 cm)



Caryopsis colour: Yellow white
Grain Shape (dorsal view): Circular
Grain Shape (profile view): Circular
Size of mark of germ: Medium
Weight of 1000 grains: Medium

Fig. 3. Important DUS characteristics of variety GDJ 1(DS 172)

Table 8. Rating of incidence of diseases at Surat and Deesa centre

i. Leaf blight (1-9)								
Trial	Year	Locations	GDJ 1	GJ 39	GJ 42	GNJ 1	GJ 43	CSV 20
LSVT	K-2018	Surat	4.3	4.7	3.7	4.3	-	4.3
LSVT	K-2019	Surat	4.3	-	-	4.3	4.7	5.0
		Mean	4.3	4.7	3.7	4.3	4.7	4.65
		Range	-	-	-	-	-	4.3-5.0
LSVT	K-2019	Deesa	3.3	-	-	5.0	3.7	4.7
		Mean	3.3	-	-	5.0	3.7	4.7
		Over all mean	4.0 (MR)	4.7	3.7	4.5	4.2	4.7
ii. Anthracnose (1-9)								
LSVT	K-2018	Surat	3.3	4.7	4.0	4.0	-	4.3
LSVT	K-2019	Surat	4.3	-	-	4.7	4.7	4.7
		Mean	3.8	4.7	4.0		4.7	4.5
		Range	3.3-4.3	-	-	4.0-4.7	-	4.3-4.7
LSVT	K-2019	Deesa	4.3	-	-	4.7	3.7	4.7
		Mean	4.3	-	-	4.7	3.7	4.7
		Over all mean	4.0 (MR)	4.7	4.0	4.5	4.2	4.6
iii. Grain mold disease score (1-9)								
LSVT	K-2018	Surat	4.0	3.3	4.0	3.7	-	4.0
LSVT	K-2019	Surat	4.3			4.7	4.3	4.7
		Mean	4.2 (MR)	3.3	4.0	4.2	4.3	4.4
iv. Ergot disease score (1-9)								
LSVT	K-2018	Surat	3.7	5.7	4.0	3.7	-	5.3
LSVT	K-2019	Surat	4.3			4.3	4.7	5.3
		Mean	4.0 (MR)	5.7	4.0	4.0	4.7	5.3

Where MR= moderately resistant

Table 9. Rating of incidence of diseases in the initial varietal trial (IVT) in the year *Kharif* 2018 at different Zones of India

i. Leaf blight (1-9)				ii. Grain mould disease score (1-9)			
Locations	GDJ 1	CSV 17	CSV 20	Locations	GDJ 1	CSV 17	CSV 20
Parbhani	3.0	2.0	2.3	Parbhani	4.7	6.0	4.7
Akola	1.0	1.0	1.0	Akola	3.7	4.0	3.0
Dharwad	1.3	1.7	1.0	Dharwad	2.7	4.0	3.3
Range	1.0-3.0	1.0-2.0	1.0-2.3	Range	2.7-4.7	4.0-6.0	3.0-4.7
Mean	1.8 (R)	1.6	1.4	Mean	3.7 (MR)	4.7	3.7
iii. Anthracnose (1-9)				iv. Ergot disease score (1-9)			
Pantnagar	5.5	6.5	5.8	Dharwad	2.0	3.0	8.0
Mean	5.5	6.5	5.8	Mean	2.0 (R)	3.0	8.0

Where R= Resistant; MR= moderately resistant

DNA fingerprinting of variety GDJ 1 along with four checks (GJ 43, GJ 39, GNJ 1 and CSV 20) was performed using 11 ISSR primers. Out of 11 primers, 7 primers (ISSR 2, ISSR 3, ISSR 7, ISSR 8, ISSR 22, ISSR 23 and ISSR 25) were showed polymorphic bands between DS-172 and checks used in fingerprinting. Polymorphic bands were demonstrated using the arrow symbol in Fig. 4.

The variety GDJ 1 (Banas Surya) found superior against checks GJ 39, GJ 42, GJ 43, GNJ 1 and CSV 20 for grain yield with an average increment of 50.0, 50.7, 19.6, 21.5 and 41.8 per cent, respectively over the years and locations. It was also found superior against respective checks for dry fodder yield with an average increment of 49.8, 52.3, 10.9, 49.6 and 34.2 per cent, respectively over the years

Table 10. Rating of incidence of Insect-pests at Surat and Deesa centre

i. Shoot fly dead heart (%)								
Trial	Year	Locations	GDJ 1	GJ 39	GJ 42	GNJ 1	GJ 43	CSV 20
LSVT	K-2018	Surat	38.12	30.59	31.31	28.84	-	36.03
		Mean	38.12	30.59	31.31	28.84	-	36.03
LSVT	K-2019	Deesa	26.82	-	-	28.69	32.13	41.40
LSVT	K-2019	Surat	22.07	-	-	22.35	29.90	31.56
		Mean	24.45			25.52	31.02	36.48
		Range	22.07-26.82	-	-	22.35-28.69	29.90-32.13	31.56-41.40
		Over all mean	29.00	30.59	31.31	26.31	31.02	36.33
ii. Stem borer dead heart (%)								
LSVT	K-2018	Surat	37.14	34.74	29.73	32.51	-	36.65
LSVT	K-2019	Surat	25.10	-	-	27.20	29.50	36.33
		Mean	31.12	34.74	29.73	29.86	29.50	36.49
		Range	25.10-37.14	-	-	27.20-32.51	-	36.33-36.65
LSVT	K-2019	Deesa	31.57	-	-	36.61	34.23	44.47
		Mean	31.57	-	-	36.61	34.23	44.47
		Over all mean	31.27	34.74	29.73	31.11	31.87	39.15




Fig. 4. DNA profiling of variety GDJ 1 with Checks.

Table 11. Rating of incidence of insect-pests in the initial varietal trial (IVT) in the year Kharif 2018 at different Zones of India

Locations	Shoot fly dead heart (%)			Stem borer dead heart (%)		
	GDJ 1	CSV 17	CSV 20	DS 172	CSV 17	CSV 20
Akola	67.9	57.6	65.7	3.5	3.7	3.6
Coimbatore	22.2	26.1	20.5	20.6	39.5	40.0
Dharwad	55.5	55.5	57.2	62.2	55.5	57.2
Indore	63.1	68.3	59.0	26.2	50.6	36.1
Palem	58.0	46.9	54.7	17.6	17.4	20.3
Parbhani	77.9	60.1	68.8	3.9	0.0	0.0
Rahuri	95.5	93.6	91.9	4.5	5.0	0.0
Surat	37.9	24.9	63.9	31.2	20.2	15.3
Udaipur	36.4	24.9	29.7	4.3	5.9	5.3
Range	22.2-95.5	24.9-93.6	20.5-91.9	3.5-62.2	0.0-55.5	0.0-57.2
Mean	57.2	50.9	56.8	19.3	21.9	19.7

and locations in the north Gujarat semi-arid zone. Further, this genotype has desirable characteristics like the tall stature of the plant with very long and broad leaves. The long symmetric semi-loose panicle has a medium neck with long branches and medium glume length. The grain is medium bold and yellow-white. It is moderately resistant to diseases like leaf blight, anthracnose, grain mold, ergot and lower infestation of shoot fly and stem borer than checks. It exhibited more crude protein in both grain (10.10 %) and dry fodder (9.20%), grain having low tannin (0.215 mg/g), while the rest of quality traits more or less similar in compare to check.

REFERENCES

- Amigot, S. L., Basílico, J. C., Bottai, H., Basílico, M. L. Z. and Fulgueira, C. L. 2005. Estrategias para mejorar la calidad de forrajes. × Congreso Argentino de Micología y XX Jornadas Argentinas de Micología. (Buenos Aires-Argentina, 22-25/05/2005) pp: 125
- Anonymous, 2017-2018. Livestock Production Statistics of India – 2017. Web: <https://www.vetextension.com/current-livestock-animal-husbandry-statistics-india/>
- Anonymous, 2018-2019. AICRP sorghum progress report; Kharif sorghum 2018.
- Singh, U. C., Mirsa, U. S., Dhamdhare, S. V. and Dwivedi, U. S. 1985. Carryover of stalk borer *Chilo partellus* (Swinhoe.) in offseason in different crops. *Journal of Entomological Research*, **9**. 170- 173
- Thakur, R. P., Reddy, B. V. S., Indira, S., Rao, V. P., Navi, S.S., Yang, X. B. and Ramesh, S. 2006. Sorghum grain mould. In: Inf. Bull. No. 72. International Crops Research Institute for the Semi-Arid Tropics, Patancheru 502324, Andhra Pradesh, India.
- Thakur, S. R., Devasthali, S. Upadhayay, S. N. and Marabi, R. S., 2019. Field Screening of Sorghum Genotypes for Resistance to Shoot fly, *Atherigona soccata* and Stem borer, *Chilo partellus*. Bulletin of Environment, Pharmacology and Life Sciences Bull. Env. Pharmacol. *Life Sci.*, **8** (5): 62-67
- Tonapi, V. A., Patil, J. V., Dayakan, B., Elengovan, M., Venkatesh, B. and Rao, K. V. R. 2011. Sorghum vision 2030. Directorate of Sorghum Research, Rajendranagar Hyderabad (AP), India. pp. 38.