



## Research Note

# Study on combining ability and gene action for yield and its attributes in eggplant (*Solanum melongena* L.)

M. A. Vaddoria and P. S. Ramani

Vegetable Research Station, Junagadh Agricultural University, Junagadh - 362 001, Gujarat  
Department of Genetics and Plant Breeding, College of Agriculture, J.A.U, Junagadh, Gujarat

(Received: 10<sup>th</sup> Jun 2015; Accepted: 31<sup>st</sup> Jul 2015)

### Abstract

Genetical study on fruit yield per plant and its attributing traits along with quality traits like total soluble solids in brinjal were conducted by following line x tester mating design comprising of seven lines and three testers at Instructional Farm, College of Agriculture, Junagadh Agricultural University, Junagadh. The analysis of combining ability revealed the predominant role of both additive and non-additive gene effects in the inheritance of yield and component traits. The genotypes NSR-1, JBG-10-208 and Pant Rituraj, displayed high gca effect for fruit yield per plant and some desirable traits like days to 50 % flowering, days to first picking, fruit girth, fruit weight, numbers of primary branches per plant, plant spread and total soluble solids. The estimates of sca effect of the crosses indicated that five hybrids manifested significant and positive sca effect for fruit yield per plant. Out of these best three specific combinations were JBG-10-208 x GJB-2 followed by ABR-2-23 x GBL-1 and AB-09-1 x GOB-1. The cross JBG-10-208 x GJB-2 was also found good specific combiner for plant height, fruit weight and fruit girth and number of primary branches per plant and also showed maximum *per se* performance. These hybrids may be exploited for obtaining transgressive segregants toward developing hybrid varieties in brinjal.

### Keywords

Brinjal, Combining ability, fruit yield

Brinjal (*Solanum melongena* L.), also known as eggplant, is an important solanaceous vegetable crop grown round the year in India. It is grown for its immature, unripe fruits which are used in the variety of ways as cooked vegetable in curries. It is grown extensively in all parts of India except at higher altitude. It is popular among people of all social strata and hence, it is rightly called as vegetable of masses. (Patel and Sarnaik, 2003). With increasing popularity of F<sub>1</sub> hybrids in brinjal, it is imperative to obtain such hybrids, having excellent quality coupled with high yields. A knowledge of general combining ability (gca) and specific combining ability (sca) helps in choice of parents or hybrids and the nature of gene action acts as a basis for choosing effective breeding methodology. The present investigation therefore was undertaken to identify potential parental combinations in order to have superior hybrids of excellent qualities coupled with high fruit yield per plant in brinjal.

The experimental material comprised of parents and their F<sub>1</sub> derived by crossing seven lines *viz.*, JBL-08-08, AB-09-1, JBG-10-208, JBL-10-04, NSR-1, JB-12-06 and ABR-2-23 and four testers *viz.*, GOB-1, GJB-2, GBL-1 and Pant Rituraj in a line x tester fashion. Eleven parental types were selected on the basis of morphological variability for growth, maturity, fruit size and shape, fruit yield and yield contributing characters. The experiment was laid out in a randomized block

<http://ejplantbreeding.com>

design with three replications during *Kharif*-2014 at Instructional Farm, College of Agriculture, Junagadh Agricultural University, Junagadh. The observations were recorded on 11 characters *viz.*, days to 50% flowering, days to first picking, fruit length (cm), fruit girth (cm), fruit weight (g), number of fruits per plant, number of primary branches per plant, plant height (cm), plant spread (cm), Total Soluble Solids and fruit yield per plant (kg). Combining ability analysis was carried out by the method suggested by Kempthorne (1957).

Analysis of variance for combining ability indicated that the mean squares due to lines were significant for all the characters. Whereas, the mean squares due to testers were significant for all the characters except for days to 50 % flowering and days to first picking. The mean squares due to lines x testers were found significant for all the characters studied indicating predominant role of both additive and non-additive gene effects in the inheritance of all these traits. (Table 1).

The ratio of  $\sigma^2_{gca}/\sigma^2_{sca}$  was less than unity for days to 50 % flowering, days to first picking, fruit weight, number of fruits per plant, number of primary branches per plant, fruit yield per plant, plant spread and total soluble solids which indicated the importance of non-additive gene action for remaining characters *viz.*, fruit length, fruit girth and plant height, the ratio of  $\sigma^2_{gca}/\sigma^2_{sca}$  was more than unity indicating the

importance of additive gene action. These results are in accordance with those obtained by Vaddoria *et al.* (2004) and Shanmugapriya *et al.* (2009) for days to 50 % flowering, numbers of primary branches per plant and fruit yield per plant; Sao and Mehta (2010) and Pachiyappan *et al.* (2012) for days to first picking, numbers of fruits per plant and plant height; Patel *et al.* (2013) and Ansari and Singh (2014) for plant height.

The estimates of gca effect indicated that among the lines, NSR-1 ranked first as it was good general combiner for most of the characters *viz.*, days to 50 % flowering, days to first picking, fruit girth, fruit weight, number of primary branches per plant, plant spread and fruit yield per plant (Table 2). Other good general combiners identified for different characters were JBG-10-208 for four character *viz.*, fruit girth, fruit weight, total soluble solids and fruit yield per plant; Pant Rituraj for fruit girth, fruit weight and fruit yield per plant; JB-12-06 for four characters *viz.*, number of fruits per plant, number of primary branches per plant, plant height and total soluble solids. Among the testers, GBL-1 was good general combiner for two characters *viz.*, fruit length and number of primary branches per plant and least fruit borer infestation. Whereas, GJB-2 was good general combiner for three characters *viz.*, fruit length, number of fruits per plant and plant height. Good general combining ability effect was exerted for plant height and total soluble solids by GOB-1. For days to 50 % flowering, days to first picking, number of primary branches per plant and plant height, the line ABR-2-23 was exerted good general combining ability effects. In view of this, NSR-1, Pant Rituraj, JBG-10-208, ABR-2-23 and JB-12-06 offer the best possibilities of exploitation for the development of improved varieties with enhanced fruit yielding ability in brinjal and these lines turned out to be good general combiners and can be used as donor parents in fruit yield improvement programme. Similar finding has also been reported by Das and Barua (2001), Singh *et al.* (2003), Kamal *et al.* (2006) and Kumar *et al.* (2012) in brinjal.

The estimates of sca effect of the crosses indicated that five hybrids manifested significant and positive sca effect for fruit yield per plant. These best five specific combinations were JBG-10-208 x GJB-2 followed by ABR-2-23 x GBL-1, AB-09-1 x GOB-1, JBL-08-8 x GJB-2 and JB-12-06 x GOB-1 (Table 3). The cross JBG-10-208 x GJB-2 was also found good specific combiner for plant height, fruit weight and fruit girth and number of primary branches per plant. Likewise, the cross ABR-2-23 x GBL-1 also showed desirable sca effect for number of fruits per plant and number of primary branches per plant and AB-09-1 x GOB-1 showed desirable

sca effect for fruit length plant height and least fruit borer infestation. The high sca effect observed for fruit yield per plant was associated with its component characters like plant height, fruit girth, fruit length, fruit weight, number of fruits per plant and number of primary branches per plant. Hybrid, JBL-08-8 x GJB-2 recorded significant sca effects in desirable direction for fruit yield per plant and fruit length. Thus, these hybrids can be exploited in practical plant breeding for selection of better transgressive segregants and they may also be exploited through heterosis breeding programme in brinjal.

#### References

- Ansari, M. J. and Singh, Y. V. 2014. Combining ability effects for fruit characters in brinjal (*Solanum melongena* L.). *Electronic J. Pl. Breeding.*, **5** (3): 385-393
- Das, G. and Barua, S. N. 2001. Heterosis and combining ability for yield and its component in brinjal. *Ann. Agric. Res. New Series.*, **22**(3): 399-403.
- Kamal, D., Bal, S. S., Kumar, A. and Sidhu, A. S. 2006. Heterosis and combining ability studies in Brinjal (*Solanum melongena* L.). *Haryana J. Hort. Sci.*, **35**(1&2): 161-165.
- Kumar, S. R., Arumugam, T., Anandakumar, C. R. and Rajavel, D. S. 2012. Estimation of heterosis and specific combining ability for yield, quality, pest and disease incidence in eggplant (*Solanum melongena* L.). *Bull. Env. Pharmacol. Life Sci.*, **2** (1): 3-15
- Kemphorne, O. 1957. An Introduction to Genetic Statistics. John Willey and Sons. Inc., New York. pp. 468-470.
- Pachiyappan, R. K.; Saravanan and Kumar, R. (2012). Combining ability analysis in eggplant (*Solanum melongena* L.). *Golden Res. Thoughts.*, **2**(2): 2-5.
- Patel, A. P., Ravindrababu, Y., Patel, A. M and Prajapati, D. B. 2013. Combining ability studies in brinjal (*Solanum melongena* L.). *GAU Research Journal.*, **38** (1): 13-16.
- Patel, K.K. and D.A. Sarnaik, 2003. Performance study of long fruited genotypes of brinjal under Raipur conditions. *The Orissa J. Hort.*, **31**(1): 74-77.
- Sao, A. and Mehta, N. 2010. Heterosis in relation to combining ability for yield and quality attributes in brinjal (*Solanum melongena* L.). *Electronic J. Pl. Breeding.*, **1** (4): 783-788
- Shanmugapriya, P., Ramya, K. and Senthikumar, N. 2009. Studies on combining ability and heterosis for yield and growth parameters in brinjal (*Solanum melongena* L.). *Crop Improvement.*, **36**(1): 68-72
- Vaddoria, M. A., Mehta, D. R., Bhatiya, V. J. and Dobariya, K. L. 2004. Combining ability for earliness and plant stature in brinjal (*Solanum melongena* L.). *Gujarat J. Applied Hort.*, **4**(1): 71-77



Table 1. Analysis of variance for combining ability and variance components for different characters in brinjal

Source	df	Days to 50 % flowering	Days to 1 <sup>st</sup> picking	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)	Number of fruits per plant
Replications	2	20.76*	10.48	0.79	0.03	23.22	4.28
Lines	6	27.19**	27.10**	3.82**	3.04**++	3133.30**++	82.90**+
Testers	3	9.63	4.48	18.73**++	5.36**++	1342.93**	24.64**
Lines x Testers	18	30.81**	28.62**	1.91**	0.67**	519.33**	27.58**
Error	54	4.47	4.21	0.56	0.17	33.19	5.09
Variance Components							
$\sigma^2_{gca}$		0.84	0.70	0.64	0.24	133.63	2.95
$\sigma^2_{sca}$		8.71	8.13	0.45	0.16	162.04	7.49
$\sigma^2_{gca}/\sigma^2_{sca}$		0.09	0.08	1.44	1.46	0.82	0.39

  

Source	df	Number primary of branches per plant	Plant height (cm)	Plant spread (cm)	TSS	Fruit yield per plant (kg)
Replications	2	0.08	12.06	19.66	0.022	0.006
Lines	6	1.77**	583.31**++	558.88**	0.738**	0.3085**
Testers	3	2.77**	116.94**	140.48**	0.621**	0.1016*
Lines x Testers	18	1.44**	68.91**	169.51**	1.104**	0.1812**
Error	54	0.17	14.28	31.03	0.054	0.026
Variance components						
$\sigma^2_{gca}$		0.12	20.35	19.31	0.037	0.010
$\sigma^2_{sca}$		0.42	18.21	46.16	0.350	0.051
$\sigma^2_{gca}/\sigma^2_{sca}$		0.30	1.11	0.41	0.108	0.209

\*, \*\* Significant at 5 and 1 per cent levels, respectively

+, ++ Significant at 5 and 1 per cent levels, respectively against lines x testers interaction.



Table 2. General combining ability effects for different characters in brinjal

Sr. No.	Parents	Days to 50 % flowering	Days to first picking	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)	Number of fruits per plant	Number of primary branches per plant	Plant height (cm)	Plant spread (cm)	Total Soluble Solids (°B)	Fruit yield per plant (kg)
Lines												
1	JBL-08-08	1.83 **	1.69 **	0.63 **	-0.62 **	-18.54**	3.09 **	-0.36 **	-0.58	-9.67 **	-0.26 **	-0.12 **
2	AB-09-1	-1.17	-1.31 *	0.58 **	-0.14	-4.90 **	-0.20	-0.27 *	0.64	-3.10	-0.36 **	-0.13 **
3	JBG-10-208	1.50 *	1.44 *	-0.28	0.66 **	13.67 **	-2.09 **	-0.39 **	-0.33	2.28	0.16 *	0.15 **
4	JBL-10-04	0.67	0.94	0.48 *	-0.40 **	3.08	-1.95 **	-0.17	-1.73	-3.14	0.11	-0.16 **
5	NSR-1	-1.75 **	-1.23 *	-0.59 **	0.69 **	25.92 **	-2.74 **	0.54 **	-12.88 **	12.68 **	-0.11	0.27 **
6	JB-12-06	0.58	0.52	-0.71 **	0.02	-18.16**	4.02 **	0.25 *	9.80 **	0.40	0.33 **	-0.01
7	ABR-2-23	-1.67 **	-2.06 **	-0.12	-0.21	-1.07	-0.12	0.39 **	5.08 **	0.55	0.13	-0.01
	SE(g <sub>i</sub> )	0.61	0.59	0.21	0.12	1.66	0.65	0.12	1.09	1.60	0.07	0.05
Testers												
1	GOB-1	-0.65	-0.15	-0.16	0.13	-3.49 **	-0.16	-0.30 **	3.53 **	1.16	0.21 **	-0.03
2	GJB-2	-0.32	-0.54	0.56 **	-0.44 **	-6.07 **	1.45 **	0.01	-1.10	1.13	0.07	-0.02
3	GBL-1	0.92	0.56	0.86 **	-0.35 **	-2.19	-0.12	0.51 **	-1.44	-3.87 **	-0.13 *	-0.05
4	Pant Rituraj	0.06	0.13	-1.26 **	0.66 **	11.75 **	-1.17 *	-0.22 *	-0.98	1.58	-0.15 **	0.10 **
	SE(g <sub>j</sub> )	0.47	0.44	0.16	0.09	1.25	0.50	0.09	0.82	1.21	0.05	0.03

\*, \*\* Significant at 5 % and 1 % levels, respectively.



Table 3. Specific combining ability effects for different characters in brinjal

Sr. No.	Crosses	Days to 50 % flowering	Days to first picking	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)	Number of fruits per plant	Number of primary branches per plant	Plant height (cm)	Plant spread (cm)	Total Soluble Solids (°B)	Fruit yield per plant (kg)
1	JBL-08-8 x GOB-1	-1.26	-2.26	-0.59	0.16	1.76	-1.61	0.12	1.52	2.12	-0.83 **	-0.14
2	JBL-08-8 x GJB-2	2.40	3.12 **	1.14 *	0.21	5.51	1.16	-0.36	0.80	4.47	-0.49 **	0.22 *
3	JBL-08-8 x GBL-1	-3.17 *	-1.31	-1.00 *	0.15	-2.13	2.91 *	0.48	-1.15	2.52	0.98 **	0.16
4	JBL-08-8 x Pant Rituraj	2.02	0.45	0.45	-0.52 *	-5.14	-2.46	-0.23	-1.17	-9.12 **	0.34 *	-0.23 *
5	AB-09-1 x GOB-1	2.40	2.40 *	0.97 *	-0.48	-8.09 *	3.09 *	-0.07	4.46 *	-1.03	-0.03	0.23 *
6	AB-09-1 x GJB-2	2.40	2.12	-0.62	-0.10	-9.91 **	2.29	0.61 *	3.33	6.40 *	0.47 **	-0.11
7	AB-09-1 x GBL-1	-5.17 **	-4.31 **	0.41	0.31	12.75 **	-4.69 **	-0.63 *	-4.56 *	-2.81	-0.13	-0.19 *
8	AB-09-1 x Pant Rituraj	0.36	-0.21	-0.75	0.27	5.25	-0.69	0.09	-3.23	-2.55	-0.31 *	0.08
9	JBG-10-208 x GOB-1	-3.60**	-1.68	-0.77	-0.02	-5.07	2.38	0.43	0.83	7.09 *	0.52 **	0.16
10	JBG-10-208 x GJB-2	1.40	0.04	-0.25	0.98 **	15.10 **	-0.78	0.82 **	4.70 *	-1.65	-0.45 **	0.37 **
11	JBG-10-208 x GBL-1	2.83 *	1.94	1.24 **	-0.77 **	-22.52 **	-0.31	-0.80 **	-4.44 *	-11.06**	0.35 **	-0.51 **
12	JBG-10-208 x Pant Rituraj	-0.64	-0.30	-0.21	-0.20	12.49 **	-1.29	-0.45	-1.09	5.62	-0.42 **	-0.02
13	JBL-10-04 x GOB-1	3.57 **	4.15 **	-0.35	-0.13	-9.87 **	-0.61	0.77 **	-7.20 **	-5.08	0.70 **	-0.22 *
14	JBL-10-04 x GJB-2	-2.10	-3.80 **	0.12	-0.71 **	-18.35 **	3.78 **	-0.06	-4.68 *	-0.37	-0.03	-0.01



Contd.....

Sr. No.	Crosses	Days to 50 % flowering	Days to 1st picking	Fruit length (cm)	Fruit girth (cm)	Fruit weight (g)	Number of fruits per plant	Number of primary branches per plant	Plant height (cm)	Plant spread (cm)	Total Soluble Solids (°B)	Fruit yield per plant (kg)
15	JBL-10-04 x GBL-1	2.00	1.11	0.32	0.65 **	15.09 **	-1.54	-0.14	10.47 **	4.69	-0.80 **	0.15
16	JBL-10-04 x Pant Rituraj	-3.48 **	-1.46	-0.09	0.20	13.13 **	-1.63	-0.57 *	1.41	0.76	0.13	0.07
17	NSR-1 x GOB-1	-1.01	-1.68	0.73	0.01	3.49	-1.35	-0.60 *	0.92	-6.15	-0.05	-0.15
18	NSR-1 x GJB-2	-0.68	-1.63	-0.31	-0.25	8.10 *	-1.37	-0.53 *	2.00	6.34	-0.68 **	-0.05
19	NSR-1 x GBL-1	-1.58	-1.39	-1.00 *	0.26	0.70	0.78	0.96 **	0.49	3.63	0.69 **	0.14
20	NSR-1 x Pant Rituraj	3.27 **	4.70 **	0.57	-0.02	-12.29 **	1.94	0.17	-3.41	-3.81	0.04	0.07
21	JB-12-06 x GOB-1	1.32	0.24	-0.54	-0.03	0.46	1.35	-0.59 *	-2.07	-3.12	-0.08	0.19 *
22	JB-12-06 x GJB-2	-4.01 **	-3.38 **	-0.04	0.15	10.01 **	-4.34 **	0.21	-3.54	-11.63 **	0.42 **	-0.15
23	JB-12-06 x GBL-1	5.08 **	4.86 **	0.84	-0.19	2.30	-1.53	-0.71 **	-2.44	-0.63	-0.41 **	-0.11
24	JB-12-06 x Pant Rituraj	-2.39	-1.71	-0.27	0.07	-12.77 **	4.52 **	1.08 **	8.05 **	15.38 **	0.08	0.08
25	ABR-2-23 x GOB-1	-1.43	-1.18	0.55	0.49 *	17.32 **	-3.25 *	-0.04	1.54	6.18	-0.22	-0.06
26	ABR-2-23 x GJB-2	0.57	3.54 **	-0.04	-0.28	-10.46 **	-0.74	-0.69 **	-2.61	-3.56	0.75 **	-0.27 **
27	ABR-2-23 x GBL-1	-0.01	-0.89	-0.81	-0.41	-6.19	4.38 **	0.83 **	1.62	3.67	-0.68 **	0.36 **
28	ABR-2-23 x Pant Rituraj	0.86	-1.46	0.30	0.20	-0.67	-0.39	-0.09	-0.56	-6.28	0.14	-0.04
	SE (S <sub>ij</sub> )	1.22	1.18	0.43	0.24	3.32	1.30	0.24	2.18	3.21	0.13	0.09