



# Electronic Journal of Plant Breeding

## Research Note

### Study of genetic variability and heritability in Indian bean (*Lablab purpureus* L.)

**U. C. Gamit, L. L. Jivani\*, H. P. Ponkia, A. Balas and A. T. Vadavia**

\*Vegetable Research Station, Junagadh Agricultural University, Junagadh- 362001, Gujarat, India  
E-Mail: lljivani@jau.in/lljivani@gmail.com

#### Abstract

The investigation was carried out at Vegetable Research Station, Junagadh Agricultural University, Junagadh (Gujarat) during late Kharif 2017-18 in Randomized Block Design with three replications to assess genetic variability in Indian bean comprised of 36 genotypes and 4 check varieties. The analysis of variance revealed that the mean sum of squares due to genotypes were highly significant for all the twelve characters. High phenotypic and genotypic coefficients of variations and high heritability coupled with high genetic advance as per cent of mean were observed for the number of primary branches per plant, the number of pods per plant, pod length, 10- green pod weight, green pod yield per plant and length of twig.

#### Keywords

Genetic variability, heritability, GCV, PCV, Indian bean

*Lablab purpureus*L. (Syn. *Dolichos lablab* L., 2n=22) is one of the important legumes as well as vegetable crops cultivated in the tropical region of Asia, Africa and America. As most of the species of Indian bean are endemic to Africa and only a few are native in India, most probably Africa is the main centre and India is the secondary centre of origin for this crop (Dana, 1976). This is one of the excellent pod vegetable crops with multipurpose used grown in India. The green pods and tender leaves are popular vegetables. It is one of the major sources of protein in the dietary of the working class, especially of whole Gujarat. It is commonly called Hyacinth bean, Bonavist bean, Indian bean, Field bean, Egyptian bean, 'Walpapadi or Valor' in Gujarat state. It has anti-diabetic property and is good for natural cure of bladder burns and cardiac problems, diarrhoea, sciatica and tenesmus. Being a legume, it can fix atmospheric nitrogen in the soil to the extent of 170 kg/ha. To develop high yielding varieties for a systematic breeding programme, information on genetic variability is basic pre-requisite. The success of any breeding programme depends upon the amount of genetic variability present in the available germplasm of a particular crop. Wider the genetic variability, more the chances of improvement through selection. Pod yield and seed yield are governed by polygenic genes control and are highly influenced by the fluctuations in the environments. Hence, the selection of plants based directly on pod yield

would not be very much reliable in many cases. Therefore, the present study was conducted to estimate genetic variability, heritability and genetic advance for pod yield and yield contributing characters.

The present investigation was carried out to assess the genetic variability in Indian bean (*Lablab purpureus* L.). The experiment was conducted at Vegetable Research Station, Junagadh Agricultural University, Junagadh (Gujarat) during late Kharif 2017-18. The 36 genotypes + four check varieties of Indian bean were raised in randomized block design with three replications. Five randomly selected plants from each genotype were considered for observation on different characters viz., days to first flowering, days to 50% flowering, days to first picking, days to the last picking, the number of pickings, twig length (cm), the number of primary branches per plant, the number of pods per plant, pod length (cm), pod width (cm), 10- green pod weight (g) and green pod yield per plant (g). The analysis of variance for Randomized Block Design (RBD) was done for each character as per Panse and Sukhatme (1985). Phenotypic Coefficient Variation (PCV) and Genotypic Coefficient Variation (GCV) were calculated as per the formula suggested by Burton and De Vane (1952). Heritability and the genetic advance was estimated using the formula suggested by Allard (1960). The analysis was done using Indostat software.

Results of analysis of variance for green pod yield and its component traits indicated (**Table 1**) that mean squares due to genotypes were found to be significant for all the traits and indicated the presence of sufficient amount of variability among the genotype for green pod

yield and its component characters in Indian bean. This result was in accordance with reports of several workers like Pan *et al.* 2001; Hotti, 2010; Dewangan *et al.*, 2017, Hadvani, 2018; Sadak Peer, 2018.

**Table 1. Analysis of variance for twelve characters in 40 genotypes of Indian bean**

Source	d. f.	Mean squares					
		No. of branches per plant	No. of pods per plant	Pod length (cm)	Pod width (cm)	10 green pod weight (g)	Green pod yield per plant (g)
Replications	2	0.0630	2097.7443	0.7024	0.0156	1.5890	36427.0234*
Genotypes	39	17.9871**	44090.7498**	22.9708**	0.2525**	417.5826**	253396.7864**
Error	78	0.0513	1922.9515	0.4067	0.0098	10.1721	9540.4024

**Table 1. Contd.**

Source	d. f.	Mean Squares					
		Days to first flowering	Days to 50 % flowering	Days to first picking	Days to the last picking	No. of pickings	Length of twig (cm)
Replications	2	2.4508	3.9083	7.6972	22.9043	0.2810	121.6029
Genotypes	39	73.2413**	94.7658**	98.0542**	32.6382**	2.8504**	5441.6095**
Error	78	4.9815	3.4126	7.2071	9.0479	0.2038	82.5506

The value of the phenotypic coefficient of variation was higher than corresponding genotypic coefficient of variation (**Table 2**) (Hotti, 2010; Hadvani, 2018; Sadak Peer, 2018) indicating the influence of environmental factors. The number of primary branches per plant exhibited the maximum values for genotypic and phenotypic coefficients of variation (Maharnavar, 2013) followed by the number of pods per plant (Chattopadhyay

and Dutta, 2010 and Hadvani, 2018), pod length (Inamdar, 2014), 10-green pod weight, green pod yield and length of twig (Inamdar, 2014; Hadvani, 2018). Moderate values of genotypic and phenotypic variations were observed for pod width and the number of pickings. The lowest value of genotypic and phenotypic variation was observed for days to 50% flowering followed by days to first flowering, days to first picking and days to the last picking.

**Table 2. Mean, range of variation, coefficient of range, phenotypic and genotypic coefficients of variation, heritability (broad sense), genetic advance and genetic advance expressed as percentage of mean for 12 characters in 40 genotypes of Indian bean**

Characters	Mean	Range	Coefficient of range (%)	Phenotypic coefficient of variation (PCV) (%)	Genotypic coefficient of variation (GCV) (%)	Heritability (Broad sense) (%)	Genetic advance	Genetic advance expressed as per cent of mean (%)
Days to first flowering	59.39	74.73-46.40	23.38	8.31	8.03	93.20	9.48	15.97
Days to 50 % flowering	62.29	77.00-50.33	26.76	9.52	9.34	96.40	11.16	18.90
Days to first picking	89.55	106.43-81.40	13.32	6.38	6.14	92.65	10.91	12.18
Days to last picking	138.45	143.00-127.60	4.81	2.38	2.02	72.28	4.91	3.54
No. of pickings	6.49	8.27-3.73	38.11	15.00	14.43	92.85	1.86	28.70
Length of twig(cm)	162.17	263.87-60.40	62.74	26.26	26.06	98.48	86.40	53.27
No. of primary branches per plant	4.13	17.87-2.33	74.62	59.25	59.17	99.71	5.02	121.71
No. of pods per plant	305.50	629.47-98.93	72.83	39.68	38.80	95.64	238.84	78.17
Pod length (cm)	8.43	15.81-4.39	56.59	32.81	32.52	98.23	5.59	66.40
Pod width (cm)	1.78	2.35-1.21	31.42	16.27	15.95	96.10	0.57	32.21
10- green pod weight (g)	39.41	65.10-19.65	53.64	30.71	30.33	97.56	23.71	61.73
Green pod yield per plant (g)	1083.02	1643.67-410.00	60.07	26.84	26.33	96.23	576.15	53.19

In the present study, high heritability (> 60%) was observed for all the 12 characters studied. The maximum heritability estimates were observed for the number of primary branches per plant (Magalingam *et al.*, 2013) followed by twig length, pod length (Hadvani, 2018), 10 green pod weight (Inamdar, 2014), days to 50% flowering (Dewangan *et al.*, 2017), green pod yield per plant (Inamdar, 2014 and Verma *et al.*, 2014), pod width (Sadak Peer *et al.*, 2018), the number of pods per plant (Verma *et al.*, 2014 and Hadvani, 2018), days to first flowering (Venkatesha *et al.*, 2016 and Sadak Peer *et al.*, 2018), the number of pickings, days to first picking (Chaitanya *et al.*, 2014) and days to last picking (Hadvani, 2018). Genetic advance expressed as per cent of mean was the highest for the number of primary branches per plant (Chaitanya *et al.*, 2014) followed by the number of pods per plant (Chattopadhyay and Dutta, 2010), pod length (Sadak Peer *et al.* (2018)), 10 green pod weight (Patel, 2014), twig length

(Chaitanya *et al.* 2014), green pod yield per plant (Hadvani, 2018), pod width and the number of pickings. The moderate magnitude of genetic advance expressed as per cent of mean was observed for characters like; days to 50% flowering, days to first flowering, and days to first picking. Whereas, days to the last picking had a low magnitude of genetic advance expressed as per cent of mean.

It can be concluded from the present investigation that high heritability coupled with high genetic advance expressed as per cent of mean was exhibited by the number of primary branches per plant, the number of pods per plant, pod length, twig length, 10- green pod weight and green pod yield per plant suggesting the existence of sufficient amount of heritable variation and wider scope for effective selection. Therefore, obtaining higher-yielding lines, the maximum weightage should be given to these attributes while making the selection.

## REFERENCES

Allard, R. W. 1960. *Principles of Plant Breeding*. John Wiley and Sons, New York.

Burton, G. W. and De Vane, E. H. 1952. Estimating heritability in tall Fescues (*Festuca arundinaceae*) from replicated clonal material. *Agron. J.*, **45**: 478-481.

Chaitanya, V.; Reddy, R. V. S. K. and Kumar, A. P. 2014. Variability, heritability and genetic advance in indigenous Dolichos bean (*Dolichos lablab* L. var. *typicus*) genotype. *Plant Archive*, **14**(1): 503-506.

Chattopadhyay, A. and Dutta, S. 2010. Characterization and identification of selection indices of pole type Dolichos bean. *73*: 33-45.

Dana, S. 1976. Origin evolution and distribution of some grain legumes. *Indian J. Genet. and Pl. Breed.*, **36**(1): 143-145.

Dewangan, R.; Bahadur, V.; Choyal P.; Ramesh, Xaxa, S.; Singh, V.P.; Sachan, S., Kerketta, A. 2017. Study on genetic variability, heritability and genetic advance in Dolichos bean genotypes (*Lablab purpureus* L.). *Int. J. Curr. Microbiol. App. Sci.*, **6**(8): 3228-3232.

Hadvani, J. K. 2018. Genetic variability, correlation, path coefficient analysis and selection indices in Indian bean. M. Sc. (Agri.) thesis. Junagadh Agricultural University, Junagadh, Gujarat.

Hotti, A. H. 2010. Genetic variability studies in field bean (*Lablab purpureus* L. Sweet). M. Sc. (Agri.) Thesis. University of Agricultural Sciences, Dharwad.

Inamdar, A. F. 2014. Genetic variability studies in pole type Dolichos bean (*Lablab purpureus*). M. Sc. (Horti.) Thesis. Mahatma Phule Krishi Vidyapeeth, Rahuri, Maharashtra.

Magalingam, V.; Yassin, M. and Ramesh Kumar, S. 2013. Genetic variability and character association in Dolichos bean. *SAARC J. Agri.*, **11**(2): 161-171.

Maharnavar, S. P. 2013. Genetic diversity and path analysis in wal (*Lablab purpureus* L.). M. Sc. (Agri.) Thesis. Mahatma Phule Krishi Vidyapeeth, Rahuri, Maharashtra.

Pan, R. S.; Singh, A. K. and Rai, M. 2001. Genetic variability in Dolichos bean. *Indian J. Plant Genet. Resour.*, **14**: 220-222.

Panse, V. G. and Sukhatme, P. V. 1985. *Statistical Methods for Agricultural Workers*. I.C.A.R., New Delhi.

Patel, S. 2014. Variability, association and genetic divergence analysis in Dolichos bean (*Lablab purpureus* L.). M. Sc. Thesis, Indira Gandhi Krishi Vishwa Vidyalaya, Raipur (C.G.).

Verma, A. K.; Jyothi, K. U. and Dorajee Rao, A. V. D. (2014). Genetic variability, heritability and genetic advances in Dolichos bean (*Lablab purpureus* L.) genotypes. *Electron. J. Plant Breed.*, **5**(2): 272-276.

Venkatesha, S. C.; Ajay, B. C.; Ramanjini Gowda, P. H.; Byregowda, M. and Ganapathy, K. N. (2016). Genetic appraisal of  $F_2$  generation of Dolichos bean for yield and attributing traits. *Electron. J. Plant Breed.*, **7**(3): 589-594.

Sadak Peer, S. Reddy, P. S. S.; Sadarunnisa, S.; Reddy, D. S. and Pandarvada, S. R. 2018. Genetic variability and heritability for yield and yield attributes in field bean (*Lablab purpureus* L.) genotypes. *Int. J. Curr. Microbiol. App. Sci.*, **7**(3): 2131-2137.