



Research Article

Assessment of combining ability effects using quality protein maize donors as testers for yield and yield traits in maize

Preeti Sharma^{2*}, Mehar Chand Kamboj² and M. S. Punia¹

¹Department of Genetics and Plant Breeding, CCS Haryana Agricultural University, Hisar- 125004, Haryana, India

²CCS Haryana Agricultural University, Regional Research Station, Karnal -132001, Haryana, India

*E-Mail: sharmapreeti.genetics@gmail.com

(Received: 26 Sep 2019; Revised: 22 Oct 2019; Accepted: 03 Nov 2019)

Abstract

The present study was accomplished to assess the general combining ability effects of parents and specific combining ability effects of hybrids for yield and yield related traits in maize. Sixty three F1 hybrids were developed and by crossing nine productive maize inbred lines with seven quality protein maize inbred testers in Line x Tester mating design and evaluated for twelve yield and yield related characters at CCSHAU, Regional Research Station, Karnal, Haryana. The combining ability analysis revealed the presence of higher magnitude of SCA than GCA variance for all characters under study except for number of grains per cob. The preponderance ratio of *sca/gca* variance revealed presence of non additive gene action in the expression of all the characters under study. Among the sixteen parents inbred lines viz., HKI 1040-4, HKI 323, HKI 161 and HKI 163 were found to be the best parent for grain yield and among the hybrids, HKI 1128 x HKI 163, HKI 659-3 x HKI 194-6, HKI 1126 x HKI 161, HKI 536YN x HKI 193-1 and HKI 488 x HKI 170(1+2) exhibited highest significant and favourable *sca* effects for yield and yield attributing characters.

Key words

gca, lines, QPM donors, *sca* and testers

Introduction

Maize (2n=2x=20), also known as corn is an allogamous crop which is widely cultivated staple food crop after wheat and rice in tropical and subtropical regions throughout the world. This crop has tremendous potential for being used in diversified sectors like human food, animal and livestock feed viz., cattle, poultry and piggery both in the form of seeds and fodder. It is also an important industrial crop as it can be used for biofuel and in starch industries. The average grain yield of maize is highest as compared to other major cereal crops such as wheat and rice. Earlier, it was much used as important staple food in Africa and Central America, and now it is gaining remarkable importance in almost all other countries including India for its diversified uses. With the increasing demand for food and fodder, India is also on threshold of maize revolution (Sharma *et al.* 2017). It is therefore, inferred that improvement in yield addressed through release of single cross hybrids. The yield potential is realized in maize mainly due to success in hybrid breeding for exploitation of heterosis in the form of hybrids and synthetics. Furthermore, the increased yield caused by heterozygosity due to outcrossing has been well documented in maize. The identification of better combining lines by evaluating parents is most essential and critical for successful exploitation of heterosis. Moreover, the combining ability and effects provide imperative insight in selection of parents that could give rise to better hybrids when

they are crossed. The value of any population depends on its potential *per se* and it's combining ability in crosses. In this context, L x T analysis (Kempthorne, 1957) has widely been used for evaluation of inbred lines by crossing them with testers (Kanagarasu *et al.*, 2010, Sundararajan and Kumar, 2011, Abrha *et al.*, 2013, Elmyhum, 2013, Kambe *et al.*, 2013). Hence, the present study has been made through systematic experimentation to study the combining ability by using QPM donors as testers for yield and yield related traits in maize and to select the appropriate parental lines, hybrid combinations with useful *gca* and *sca* effects.

Materials and Methods

Nine productive maize inbred lines (referred to as female line) were crossed with seven quality protein maize inbred lines (referred to as male tester) in Line x Tester mating design to generate sixty three experimental hybrids during *kharif* and *rabi* of 2013-14 (Table 1). The field experiment consisted of sixty three experimental hybrids and parental inbred lines along with two standard checks HQPM 1 and HM 5 was conducted at research area of maize section of Chaudhary Charan Singh Haryana Agricultural University, Regional Research Station, Karnal. The experiment was laid out in Randomized Complete Block Design in a two rows of 3m length along with three replications and border rows were also planted at each replication in order to avoid border effect. The



inter-row and intra row was kept 75 cm and 20 cm respectively. All the recommended agricultural practices of CCS Haryana Agricultural University were adopted to maintain healthy crop. Ten highly competitive plants were selected and tagged in each plot and in each replication to record the observations on various yield and yield contributing traits *viz.*, days to 50 per cent tasseling, days to 50 per cent silking; days to 50 per cent maturity, plant height, ear height, number of cobs per plant, cob length, cob diameter, number of grains per cob, hundred grain weight, grain yield per plant and shelling percentage. Mean values of each observation were collected to carry out analysis of variance for each character. At first the test of significance among all the genotypes including crosses and parents were estimated and when that were found significant then Line \times Tester analysis was performed. Combining ability analysis was performed using OPSTAT software program, to partitioned treatment sum of square into sum of squares due to parents, crosses and parents vs. crosses with appropriate degree of freedom (Singh and Chaudhary, 1985). Heritability estimates in broad sense of each trait was computed as per Falconer (1989) to determine the progress under selection. The estimates of genetic advance were obtained as per Singh and Choudhary (1985) by computing the difference between the mean of the progeny of selected individuals and the base population.

Results and Discussion

The significant differences for parents vs. hybrids indicated the presence of heterosis and further indicated significant amount of genetic variability among lines and testers for all the characters (Table 2). The results revealed that different lines as well as testers showed markedly varied combining ability effects for all the characters except for number of cobs per plant (where only *sca* effects were significant for only one cross) and cob diameter (where none of the cross exhibits significant *sca* effects) which in turn revealed importance of additivity and non-additivity in the inheritance of these characters. This warranted further estimation of *gca* and *sca* effects for characters being studied. Significant differences for *gca* and *sca* effects for the characters under study were also reported by Mutimaamba *et al.*, (2010), Singh *et al.*, (2013), Elmyhum (2013) and Hemalatha *et al.*, (2014).

The data in table 3 indicated a wide range of variability for *gca* effects among the parents for different characters. Good combiners were identified on the basis of high and significant *gca* effects towards the desired direction. Perusal of the table revealed that no line was observed to be a

good combiner for all the traits, however good combiners for more than two characters identified. Among the lines, HKI 1040-4 and HKI 323 exhibited significant positive *gca* effects for most of the traits including grain yield. Among the testers, HKI 161 and HKI 163 were considered good combiners for grain yield and other traits. Such parents contribute favourable alleles in the process of synthesis of new varieties and these potential lines and testers could be efficiently utilized. Earlier similar reports of good combiner for more than three characters were also reported by Singh and Gupta (2009); Ruth *et al.*, (2010); Elmyhum (2013); Kambe *et al.*, (2013); Hemalatha *et al.*, (2014).

High and positive *sca* effect is desirable for grain yield, cob length, cob diameter, number of grains per cob, shelling percentage and 100 grain weight. Both negative and positive and significant estimates of *sca* effects were observed among the crosses for grain yield per plant (Table 4). Magnitude of *sca* effect for grain yield per plant revealed that fifteen crosses showed significant positive *sca* effects for grain yield per plant and majority of crosses were having at least one average combiner parent. A critical evaluation of the results with respect to specific combining ability effects expressed that none of the cross combinations exhibited desirable significant *sca* effects for all the characters. The crosses HKI 1128 x HKI 163, HKI 659-3 x HKI 194-6, HKI 1126 x HKI 161, HKI 536YN x HKI 193-1 and HKI 488 x HKI 170(1+2) with high positive and significant estimates of *sca* effects for more than two traits could be selected for their specific combining ability to use in maize improvement. The *sca* effects of the crosses did not show any specific trends in cross combinations between parents possessing high, medium and low *gca*. In most of the cases, the crosses those showed high *sca* effects involved at least one good general combiner. The superiority of crosses involving high \times low combiners as parents could be explained on the basis of interaction between positive alleles from good combiners and negative alleles for the poor combiners as parents. The high yield of such crosses would be non-fixable and thus could be exploited for heterosis breeding. These results are in agreement with the findings of Iqbal *et al.* (2007) and Shams *et al.* (2010), Lal *et al.* (2011), Ruswandi *et al.* (2015), Yerva *et al.* (2016) who reported high and significant level of *sca* effects in most of the crosses they studied for grain yield in maize. Earlier reports of Xingming *et al.* (2002) also suggested that good *gca* parents play positive role in high yielding crosses. Amiruzzaman *et al.* (2011) also pointed out that the *sca* effect is a result of the interaction of *gca* effects of the parents

and that it can improve or deteriorate the hybrid expression compared to the expected effect based on *gca* only. These results are supported by, Hemavathy and Balaji (2008) while in contrary, Ivy and Howlader (2000) obtained high *sca* effect for grain yield in low \times low general combining parents and revealed that crosses with high *sca* effects did not always had parents with good *gca* effects. For other yield contributing traits most of the top ranking specific combiners revealed average specific combining ability and from the results it was concluded that *gca* effects of the parents did not reflected in their *sca* effects for all the traits. Such a relationship between *gca* and *sca* indicates the importance of epistasis and crosses are expected to produce desirable transgressive segregants. The results obtained in the present study are mostly in conformity with the earlier findings of Mahto and Ganguli (2003), Malik *et al.* (2004) and Kanagarasu *et al.* (2010) for grain yield and other component characters.

Results regarding gene action, based on variance of GCA: SCA presented in Table 5 revealed that the preponderance ratio (GCA: SCA) was less than unity for all the characters except number of grains per cob, thus, indicated that non additive gene action had played more role in these characters. The SCA variances were in general higher than the GCA variances which depict that one can go for hybrid breeding programme in future with the present set of breeding material. Similar results were reported by Jabeen *et al.* (2007), Ruth *et al.* (2010), Abrha *et al.* (2013), Singh *et al.* (2013), Elmyhum (2013), Kapoor and Lata (2013), Hemalatha *et al.* (2014), Chahar *et al.* (2014). Contrarily, importance of additive gene effects was reported by Sharma *et al.* (2004) and Alammie *et al.* (2006). The percentage contribution of line \times tester interactions was found higher than testers for all the characters except for days to 50 percent tasseling and 50 percent silking. Similar findings were reported for ear length and ear diameter by Kanta *et al.* (2005).

The study impetus that on the basis of overall performance of F_1 hybrids and parental lines, it is possible to select some lines with good *gca* and yield potential and hybrids with good *sca*. Hence, the present study contently helped in identifying superior parents such as HKI 323, HKI 1040-4, HKI 161 and HKI 163 and crosses HKI 1128 \times HKI 163, HKI 659-3 \times HKI 194-6, HKI 1126 \times HKI 161, HKI 536YN \times HKI 193-1 and HKI 488 \times HKI 170(1+2) could be exploited in the production high yielding hybrids with good protein quality combination since testers are promising QPM lines. The performance of such desirable crosses need to be evaluated first at a few location and subsequently for the top one or two hybrids in

multilocational trial at farmer field to confirm their superiority

Acknowledgements

This research was financed by Department of Science and Technology by providing INSPIRE Fellowship to the first author. All authors are indebted to the funding agency and also to the Regional Research Station for providing necessary support to conduct this research.

References

Abrha S.W., Habtamu Z.Z. and Dagne W.G., 2013. Line \times tester analysis of maize inbred lines for grain yield and yield related traits. Asian J. Plant Sci. Res., **3** (5):12-19.

Alammie A., Wali M.C., Salimath PM. and Jagadeesha B.C. 2006. Combining ability and heterosis for grain yield and ear characters in maize, Karnataka J. Agric. Sci., **19**: 13-16.

Amiruzzaman M., Islam MA., Hasan L., Kadir M. and Rohman M. 2011. Heterosis and combining ability in a diallele among elite inbred lines of maize (*Zea mays* L.). Emirates J. Food and Agric., **23**: 204-208.

Chahar S., Vyas M., Ranwah B.R and Vadodariya G., 2014. Heterosis and Combining Ability Analysis for Yield and its Contributing Traits in Early Maturing Maize (*Zea mays* L.) Genotypes. Trends in Biosci., **7** (14): 1774-1777.

Elmyhum M. 2013. Estimation of combining ability and heterosis of quality protein maize inbred lines. Afric. J. Agric. Res., **8**(48): 6309-6317.

Falconer D.S. 1989. Introduction to quantitative genetics (3rd ed.) Longman Scientific and Technical, Longman House, Burnt Mill, Harlow, Essex, England.

Hemalatha V., Sai K.R., Swarnalatha V. and Suresh J. 2014. Combining ability and gene action for morphological parameters in quality protein maize (*Zea Mays* L.). Int. J. Pl. Ani. & Environ. Sci., **4** (2): 230-235.

Hemavathy A.T. and Balaji K. 2008. Analysis of combining ability and heterosis groups of white grain quality protein maize (QPM) inbreds. Crop Res., **36** (1/3): 224-234.

Iqbal A.M., Nehvi F.A., Wani S.A., Qadir R. and Dar D.A., 2007. Combining ability analysis for yield and yield related traits in maize (*Zea mays* L.). Int. J. Pl. Breed. & Genet., **1**(2):101-105.

Ivy N.A. and Howlader M.S. 2000. Combining ability in maize. Bangladesh J. Agricul Res., **25**: 385-392.



Jabeen F., Sahib H. and Satyanarayana E., 2007. Combining ability analysis in quality protein maize (*Zea mays* L.). *Crop Res.*, **34** (1,2 & 3): 171-175.

Kambe G.R., Kage U., Kumar H.C., Lohithaswa B.G., Shekara and Hobha D. 2013. Combining ability studies in maize (*Zea mays* L.). *Mol. Pl. Breed.*, **4**(14): 116-127.

Kanagarasu S., Nallathambi G. and Ganesan K.N. 2010. Combining ability analysis for yield and its component traits in maize (*Zea mays* L.). *Elect. J. Pl. Breed.* **1**(4): 915-920.

Kanta G., Singh H.B., Sharma J.K. and Guleri G.K. 2005. Heterosis and combining ability studies for yield and its related traits in maize. *Crop Res.*, **30**: 221-226.

Kapoor C. and Lata S. 2013. Screening of some maize (*Zea mays* L.) inbreds for conversion to quality protein maize lines. *Indian J. Agric. Res.*, **47** (5): 457 – 460.

Kempthorne O. 1957. *An Introduction to Genetic Statistics*. 1st Edn., John Wiley and Sons, New York.

Lal M., Singh, D. and Dass, S. 2011. General and specific combining ability studies in maize using line x tester design. *Agric. Sci. Digest.*, **31** (1): 8 – 13.

Mahto R.N. and Ganguli D.K. 2003. Combining ability analysis in inter varietal crosses of maize (*Zea mays* L.). *Mad. Agricul. J.*, **90**: 29-33.

Malik S.I., Malik H.N., Minhas N.M. and Munir M. 2004. General and Specific combining ability studies in maize. *Int. J. Agric. and Biol.*, **6**: 856-859.

Mutimaamba C., Lungu D. and Robert J.M. 2010. Combining ability analysis of quality protein maize (QPM) and non-QPM inbred lines for kernel quality and some agronomic characteristics. *Crop Res.*, **34** (1): 171-175.

Ruswandi D., Supriatna J., Makkulawu A.T., Waluyo B., Marta H., Suryadi E. and Ruswandi S. 2015. Determination of combining ability and heterosis of grain yield components for maize mutants based on Line x Tester analysis. *Asian. J. Crop. Sci.*, **7**: 19-33.

Ruth N.M., Alpha O.D., Dan M. and Kiarie N. 2010. Combining ability of early maturing quality protein maize inbred lines adapted to Eastern Africa. *Field Crops Res.*, **119** (2-3): 231-237.

Shams M., Choukan R., Majidi E. and Darvish F. 2010. Estimation of combining ability and gene action in maize using line x tester method under three irrigation regimes. *J. Res. in Agricul. Sci.* **6**:19-28.

Sharma P., Punia M.S., Kamboj M.C., Singh N. and Mehar Chand. 2017. Evaluation of quality protein maize crosses through line x tester analysis for grain yield and quality traits. *Agricul. Sci. Digest.*, **37** (1): 42-45.

Sharma S., Narwal R., Kumar M.S. and Dass S. 2004. Line x tester analysis in maize (*Zea mays* L.), *Forage Res.*, **30**: 28-30.

Singh P.K., Singh N., Singh A.K., Shahi J.P. and Rao M. 2013. Heterosis in relation to combining ability in Quality protein maize (*Zea mays* L.). *Biolife*. **1**(2):65-69. (2013).

Singh R.K. and Chaudhary B.D. 1985. *Biometrical Methods in Quantitative Genetic Analysis*. Kalyani, New Delhi, India.

Singh S.B. and Gupta B.B., 2009. Heterotic expression and combining ability analysis for yield and its components in maize (*Zea mays* L.) inbreds. *Progres. Agricul.* **9** (2): 120-123.

Sundararajan R. and Kumar P.S. 2011. Studies on combining ability through line x tester analysis in maize (*Zea mays* L.). *Pl. Arch.*, **11**(1): 75-77.

Xingming F., Tan J., Chen Z. and Yang J. 2001. Combining ability and heterotic grouping of ten temperate, tropical and subtropical Quality Protein Maize. In: Proc. 8th Asian Reg. Maize Workshop Bangkok, Thailand.

Yerva S.R., Sekhar T.C., Allam C.R. and Krishnan V. 2016. Combining ability studies in maize (*Zea mays* L.) for yield and its attributing traits using Griffing's diallel approach. *Elect. J. Pl. Breed.*, **7**(4): 1046-1055.



Table 1. List of inbred lines used to develop experimental hybrids

S. No.	♀ (Normal and productive inbreds)	S. No.	♂ (Quality protein maize inbreds)
1	HKI 1128 (L1)	1	HKI 163 (T1)
2	HKI 288-2 (L2)	2	HKI 161 (T2)
3	HKI 659-3 (L3)	3	HKI 170(1+2) (T3)
4	HKI 323 (L4)	4	HKI 194-6 (T4)
5	HKI 1126 (L5)	5	HKI 193-1 (T5)
6	HKI 1105 (L6)	6	HKI 193-2 (T6)
7	HKI 536YN (L7)	7	HKI 5072-BT(1-2)-2 (T7)
8	HKI 488 (L8)		HQPM 1 (Check 1)
9	HKI 1040-4 (L9)		HM 5 (Check 2)

Table 2. Analysis of variance (MSS) of parents, hybrids and combining ability for different traits in a line x tester crosses of maize

Sources of variation	d.f	Days to 50 % taselling	Days to 50 % silking	Days to maturity	Plant height (cm)	Ear height (cm)	No. of cobs per plant	Cob length (cm)	Cob diameter (cm)	Number of grains / cob	100 grain weight (g)	Grain yield per plant (g)	Shelling percentage
Replication	2	1.987	1.549	1.447	103.085	21.444	1.392	2.941	0.17	229.99	3.734	75.541	22.781
Genotypes	78	18.92**	14.93**	24.73**	2923.57**	431.53**	3.51**	11.93**	0.47**	34148.92**	21.82**	1999.45**	21.32**
Females	8	19.92**	5.17**	25.91**	337.95**	352.87**	2.07**	5.78**	0.65**	4368.28**	25.09**	209.53**	18.36**
Males	6	14.93**	13.52**	25.42**	178.63**	176.54**	10.3**	6.05**	0.74**	4616.15**	13.94**	74.44	40.78**
Hybrids	62	15.16**	14.48**	13.63**	2007.08**	141.57**	3.09**	4.73**	0.20**	8731.36**	16.98**	939.18**	16.92**
Males Vs Females	1	7.64**	14.03**	8.67**	8.89	7.05	0.62	5.21*	0.35	296.56	0.1	8.28	1.02
Parents Vs Hybrids	1	279.26**	130.42**	715.49**	99815.5**	20993.1**	2.77	548.97**	14.67**	2059331.9**	365.09**	95596.8**	221.56**
Error	156	1.26	1.39	1.17	86.25	26.14	0.02	0.87	0.15	59.03	2.36	38.17	5.99
Line	8	60.31**	67.32**	31.14	6363.57**	185.33**	0.03**	4.03**	0.02	17537.68**	5.70**	1324.49**	24.34**
Tester	6	3.22**	2.91**	13.79	1216.19**	155.03**	0.06**	6.19**	0.13	5457.01**	28.81**	1182.40**	26.99**
LinexTester	48	5.70**	7.11**	10.69	1379.82**	132.59**	0.06**	4.67**	0.13**	7838.70**	17.38**	844.56**	14.42**
Error	124	1.21	1.34	0.95	91.23	29.47	0.02	1.06	0.18	47.13	1.91	45.18	2.76

*:**Significant at 5 and 1% levels respectively



Table 3. Estimates of gca effects for different traits in lines and testers of maize

Parents	Days to 50 % % taselling	Days to 50 % silking	Days to maturity	Plant height (cm)	Ear height (cm)	No. of cobs per plant	Cob length (cm)	Cob diameter (cm)	Number of grains / cob	100 grain weight (g)	Grain yield per plant (g)	Shelling percentage
Lines												
HKI 1128	2.772*	2.661**	1.603**	22.423**	-0.365	0.077	0.401	-0.024	-35.646**	0.546	-5.358**	-0.457
HKI 288-2	1.963*	1.947**	1.651**	25.804**	6.063**	0.039	0.387	-0.01	-14.788**	-0.915*	-7.492**	-0.186
HKI 659-3	-0.323	-0.672	-0.349	-16.624**	-3.556*	-0.018	-0.747*	0.009	7.354**	0.518	2.606	0.101
HKI 323	-2.466**	-2.339**	-0.873**	-8.72**	-3.556*	-0.028	0.087	-0.024	18.164**	-0.457	7.393**	-0.034
HKI 1126	-2.942**	-2.720**	-1.778**	-8.386**	1.397	0.001	0.311	0.052	10.831**	0.239	2.478	-0.771
HKI 1105	0.201	0.090	0.413	-18.624**	-3.032	-0.037	-0.037	0.047	-51.503**	-0.176	-13.82**	-1.718**
HKI 536YN	0.915**	0.947**	0.937**	-16.196**	1.635	-0.056	-0.642*	-0.029	11.545**	-0.386	0.219	-0.155
HKI 488	0.487	0.566	-0.594*	13.09**	0.968	0.001	0.430	-0.029	12.116**	0.065	1.353	1.244*
HKI 1040-4	-0.668*	-0.481	-1.111**	7.233*	-0.698	0.02	-0.289	0.009	41.926**	0.565	12.621**	1.976**
SE \pm GCA (Lines)	0.339	0.357	0.302	2.947	1.675	0.042	0.318	0.133	2.1187	0.4267	2.0743	0.5127
C.D (5 %)	0.666	0.701	0.592	5.777	3.284	0.082	0.624	0.261	4.153	0.836	4.066	1.005
C.D (1 %)	0.875	0.922	0.778	7.593	4.316	0.108	0.820	0.343	5.458	1.099	5.343	1.321
Testers												
HKI 163	0.386	0.397	1.032**	4.148	3.947**	0.083*	-0.337	0.071	0.524	1.155**	7.15**	0.274
HKI 161	-0.206	-0.122	0.143	8.63**	1.54	0.046	-0.119	0.063	-6.624**	1.539**	8.001**	1.099*
HKI 170(1+2)	-0.593*	-0.655*	-1.339**	-6.926**	-3.312*	-0.006	-0.137	-0.022	-19.365**	0.167	-3.149	1.209**
HKI 194-6	0.09	0.101	-0.153	-5.481*	-2.349	-0.051	0.819**	-0.118	13.153**	-0.501	0.342	-0.534
HKI 193-1	0.597*	0.286	0.254	-0.926	0.058	-0.043	0.063	0.026	-15.328**	-1.255**	-11.175**	-1.650**
HKI 193-2	0.127	0.175	0.254	7.333**	0.095	-0.029	0.363	0.045	12.487**	-0.896*	-3.036	0.092
HKI 5072-BT(1-2)-2	-0.392	-0.381	-0.19	-6.778**	0.021	0.001	-0.652*	-0.066	15.153**	-0.209	1.866	-0.49
SE \pm GCA (Testers)	0.299	0.315	0.266	2.599	1.477	0.037	0.280	0.117	1.8685	0.3763	1.8294	0.4522
C.D (5 %)	0.587	0.618	0.522	5.095	2.896	0.073	0.550	0.230	3.662	0.738	3.586	0.886
C.D (1 %)	0.772	0.813	0.686	6.697	3.806	0.096	0.723	0.303	4.813	0.969	4.713	1.165

*:**Significant at 5 and 1% levels respectively



Table 4. Estimates of *sca* effects for different traits in lines and testers of maize

Sr. No.	Crosses	Days to 50% taselling	Days to 50% silking	Days to maturity	Plant height (cm)	Ear height (cm)	No. of cobs per plant	Cob length (cm)	Cob diameter (cm)	Number of grains/cob	100 grain weight (g)	Grain yield per plant (g)	Shelling per centage
1	L1×T1	1.376	1.413	2.730**	46.947**	12.624**	0.508**	1.466	0.258	120.238**	0.511	27.729*	-1.705
2	L1×T2	0.635	0.931	0.619	-19.201*	-2.968	-0.055	-0.653	0.032	-57.947**	1.610	-11.322*	-1.877
3	L1×T3	0.598	0.265	0.101	-44.312**	-8.783*	-0.270*	1.099	0.217	59.794**	-2.958**	1.438	2.844*
4	L1×T4	0.339	0.376	-1.085	-7.423	-1.413	-0.025	0.477	-0.253	-45.392**	3.180**	-0.300	0.63
5	L1×T5	-0.069	0.19	0.508	7.354	2.18	0.034	-1.001	0.369	-24.91**	1.581	-0.469	0.035
6	L1×T6	-1.365	-1.698	-0.825	12.429	-1.524	-0.181	0.232	-0.416	28.608**	0.291	8.988	1.534
7	L1×T7	-1.713*	-1.476	-2.048*	4.206	-0.116	-0.011	-1.62	-0.205	-80.392**	-4.215**	-26.064**	-1.461
8	L2×T1	-1.148	-1.206	-1.317	-16.101*	-7.138	-0.054	-0.787	-0.09	-18.286**	-1.202	-11.317*	2.081
9	L2×T2	-0.889	-0.688	-1.095	26.751**	-5.063	-0.084	-1.572	0.117	4.862	-0.216	7.426	3.459*
10	L2×T3	1.407	0.979	0.72	-3.693	1.455	0.168	1.28	-0.264	-39.063**	-1.058	-11.581*	-1.62
11	L2×T4	1.148	1.09	2.868**	14.862	4.825	0.013	-1.309	-0.268	-25.249**	1.897	-1.402	-3.365*
12	L2×T5	0.074	-0.095	-0.54	-3.026	1.751	-0.195	0.08	0.021	58.566**	-2.658*	0.888	0.828
13	L2×T6	-0.556	0.016	-2.54**	17.714*	9.048*	0.057	0.447	0.236	19.418**	1.459	13.379*	0.016
14	L2×T7	-0.037	-0.095	1.905*	-36.508**	-4.878	0.094	1.861*	0.247	-0.249	1.779	2.607	-1.399
15	L3×T1	1.471	0.746	0.349	-14.339	-1.852	-0.063	-0.287	0.158	17.905**	2.289*	17.485**	1.988
16	L3×T2	-2.270	-2.402*	-1.862*	-30.153**	-8.111	-0.093	1.161	-0.035	0.053	-1.650	-10.242	-1.338
17	L3×T3	-0.307	-0.069	0.386	24.735**	5.407	-0.108	-0.587	-0.05	21.794**	-0.863	-2.973	-0.454
18	L3×T4	0.767	0.709	-1.466	-4.709	5.111	-0.130	0.424	-0.087	62.608**	2.784*	27.82**	3.612**
19	L3×T5	0.360	0.857	2.127**	-0.265	4.704	0.129	0.480	-0.331	-25.243**	-3.927**	-18.406**	-4.319**
20	L3×T6	-2.937**	-2.698**	-2.206**	12.810	-5.000	0.181	-0.920	0.017	-36.392**	1.726	-0.139	1.906
21	L3×T7	2.915**	2.857**	2.571**	11.921	-0.259	0.085	-0.272	0.328	-40.725**	-0.334	-13.544*	-1.396
22	L4×T1	-0.386	0.079	1.206	-41.243**	-10.328*	0.013	0.013	-0.576	18.762**	0.188	10.381	0.839
23	L4×T2	0.206	0.265	0.095	-27.058**	-6.921	0.05	1.395	-0.135	6.243	1.443	15.761**	1.181
24	L4×T3	-0.497	-0.402	0.243	-1.503	0.598	0.102	0.113	-0.150	-24.349**	0.495	-12.109*	-0.639
25	L4×T4	-0.423	-0.291	-0.275	-3.947	-1.698	-0.054	-0.476	0.113	-26.868**	-3.587**	-23.18**	-0.933
26	L4×T5	0.836	0.524	-1.016	41.164**	7.561	-0.128	-1.42	0.269	0.614	0.124	2.570	1.819
27	L4×T6	1.540	1.635	1.317	42.238**	12.19**	-0.010	-0.653	0.050	-25.201**	-1.012	-12.199*	-2.916*
28	L4×T7	-1.275	-1.860*	-1.571*	-9.651	-1.402	0.028	1.028	0.428	50.799**	2.348*	18.776**	0.649
29	L5×T1	0.090	-0.206	-0.556	15.423*	2.196	-0.283*	1.356	0.048	51.762**	-2.372*	-0.891	1.333
30	L5×T2	-0.651	-0.688	-2.333**	-7.058	1.270	-0.046	0.604	0.056	37.577**	1.930	21.225**	-0.686
31	L5×T3	-1.354	-1.354	-1.852*	4.497	3.455	-0.060	0.456	0.074	-23.349**	1.742	10.572	1.005
32	L5×T4	-0.280	0.090	0.630	-1.947	3.492	0.117	0.501	0.37	-22.868**	-3.537**	-19.763*	-0.553
33	L5×T5	-0.021	-0.095	1.222	4.831	0.751	0.110	1.023	-0.241	8.614	0.434	-0.589	0.333



34	L5×T6	1.016	0.683	-0.444	-12.762	-10.619[*]	0.095	-2.144[*]	0.141	-77.201^{**}	1.051	-19.128^{**}	-3.805^{**}
35	L5×T7	1.201	1.571	3.333^{**}	-2.984	-0.545	0.066	-1.796[*]	-0.448	25.466^{**}	0.751	8.573	2.373
36	L6×T1	-1.720[*]	-1.883[*]	-1.746[*]	2.661	5.291	0.089	0.704	-0.314	-25.905^{**}	0.056	-8.719	-2.654
37	L6×T2	-0.127	-0.164	0.476	3.513	6.698	-0.074	-0.448	-0.04	35.243^{**}	-4.318^{**}	-11.55[*]	-0.089
38	L6×T3	-0.164	0.169	-1.376	-0.598	-6.783	-0.089	-2.396^{**}	0.179	-88.349^{**}	2.690[*]	-6.160	-1.975
39	L6×T4	0.577	0.28	-0.894	2.958	-3.413	0.022	0.748	0.308	-25.201^{**}	-2.835[*]	-13.661[*]	-3.019[*]
40	L6×T5	-0.164	0.095	0.698	5.402	5.847	-0.052	1.67[*]	-0.103	43.614^{**}	1.406	13.936[*]	3.576^{**}
41	L6×T6	2.206[*]	2.206[*]	3.698^{**}	-9.190	-2.190	0.200	0.404	-0.154	14.132[*]	1.427	9.313	2.915[*]
42	L6×T7	-0.608	-0.905	-0.857	-4.746	-5.450	-0.096	-0.681	0.123	46.466^{**}	1.573	16.841[*]	1.246
43	L7×T1	0.233	0.460	-0.603	-3.767	-0.376	-0.025	-0.292	0.262	-67.286^{**}	1.216	-14.035[*]	-1.383
44	L7×T2	2.159[*]	1.979 [*]	4.619^{**}	5.751	2.032	0.078	1.59	0.237	1.862	-0.511	-6.792	0.572
45	L7×T3	2.122[*]	2.312[*]	2.434^{**}	15.640[*]	2.550	0.197	-0.625	-0.245	-0.063	-0.723	-1.963	-1.318
46	L7×T4	0.196	0.090	-0.418	-5.804	-2.079	-0.092	-0.014	0.118	21.418^{**}	0.935	9.186	1.304
47	L7×T5	-2.312^{**}	-2.429[*]	-1.492	-13.693	-7.487	0.034	0.008	-0.326	39.899^{**}	4.663^{**}	27.937^{**}	-1.903
48	L7×T6	-1.175	-1.317	-0.825	-16.286[*]	3.143	-0.114	0.442	-0.045	76.418^{**}	-0.630	19.614^{**}	3.892^{**}
49	L7×T7	-1.323	-1.095	-3.714^{**}	18.159[*]	2.217	-0.077	-1.110	-0.001	-72.249^{**}	-4.950^{**}	-33.948^{**}	-1.163
50	L8×T1	0.661	0.841	-0.841	-2.386	3.291	-0.216	-2.196[*]	0.129	-89.857^{**}	-3.152^{**}	-31.269^{**}	-1.045
51	L8×T2	1.921[*]	2.026[*]	0.714	30.132^{**}	10.032[*]	0.021	-0.415	0.203	-31.376^{**}	0.660	-7.313	-0.047
52	L8×T3	0.550	0.693	0.862	1.688	-9.116[*]	0.140	0.504	0.155	61.698^{**}	0.146	13.387[*]	-0.56
53	L8×T4	-0.042	-0.196	1.677[*]	-13.423	-8.746	0.117	-0.885	-0.215	55.18^{**}	1.553	19.876^{**}	2.029
54	L8×T5	0.217	-0.048	-1.397	-17.646[*]	-6.153	-0.09	1.304	0.440	-16.672^{**}	0.572	-0.407	-0.222
55	L8×T6	-1.413	-1.603	-0.063	-22.238^{**}	2.143	-0.038	0.537	-0.378	21.18^{**}	-2.658[*]	-6.666	-0.193
56	L8×T7	-1.894^{**}	-1.714	-0.952	23.873^{**}	8.550	0.066	1.152	-0.334	-0.153	2.879[*]	12.392[*]	0.038
57	L9×T1	-0.577	-0.444	0.778	12.804	-3.709	0.032	0.023	0.124	-7.333	2.465[*]	10.637	0.546
58	L9×T2	-0.984	-1.259	-1.333	17.323[*]	3.032	0.202	-1.662[*]	-0.435	3.481	1.077	2.806	-1.176
59	L9×T3	-2.354^{**}	-2.593^{**}	-1.519	3.545	11.217[*]	-0.079	0.156	0.084	31.889^{**}	0.529	9.389	2.718[*]
60	L9×T4	-2.380[*]	-2.148[*]	-1.037	19.434[*]	3.921	0.032	0.534	-0.087	6.370	-0.390	1.425	0.294
61	L9×T5	0.979	1.000	-0.111	-24.122^{**}	-9.153[*]	0.158	-2.144[*]	-0.098	-84.481^{**}	-2.295[*]	-25.461^{**}	-0.147
62	L9×T6	2.683^{**}	2.778^{**}	1.889[*]	-24.714^{**}	-7.190	-0.190	1.656[*]	0.550	-20.963^{**}	-1.655	-13.164[*]	-3.349[*]
63	L9×T7	2.534^{**}	2.667^{**}	1.333	-4.270	1.884	-0.153	1.438	-0.139	71.037^{**}	0.169	14.367^{**}	1.113
Standard Error		0.8991	0.9466	0.7993	7.7987	4.4326	0.1112	0.8421	0.3527	5.6055	1.1289	5.4882	1.3565
C.D (5 %)		1.762	1.855	1.567	15.285	8.688	0.218	1.651	0.691	10.987	2.213	10.757	2.659
C.D (1 %)		2.316	2.438	2.059	20.089	11.418	0.286	2.169	0.909	14.440	2.908	14.138	3.494



Table 5. Proportional contribution of lines, testers and their interactions to genetic components of variance or estimates of the variance due to GCA, SCA, dominance variance and additive variance for different traits in maize crosses

Sr. no.	Character	σ^2 gca	σ^2 sca	Ratio	σ^2 (A)		σ^2 (A) / σ^2 (D)	Percentage Contribution of			
					Additive	Dominance					
					gca : sca			Lines	Testers	Interaction	
1	Days to 50 % tasseling	1.32	1.90	0.69	5.27	7.60	0.69				
2	Days to 50 % silking	1.07	2.15	0.50	4.29	8.60	0.50	1.94	60.00	38.06	
3	Days to maturity	1.37	3.10	0.44	5.48	12.42	0.44	9.79	29.48	60.73	
4	Plant height (cm)	103.39	633.08	0.16	413.54	2532.33	0.16	5.86	40.91	50.23	
5	Ear height (cm)	25.91	44.99	0.58	103.64	179.97	0.58	10.60	16.89	72.51	
6	No. of cobs per plant	0.00	0.002	0.00	0.00	0.01	0.20	9.75	7.21	83.04	
7	Cob length (cm)	0.95	1.52	0.62	3.80	6.08	0.62	12.65	10.97	76.37	
8	Cob diameter (cm)	0.02	0.08	0.24	0.07	0.30	0.24	3.58	1.35	92.07	
9	Number of grains / cob	3329.38	3056.90	1.09	13317.52	12227.60	1.09	5.96	25.54	68.50	
10	100 grain weight (g)	503.14	6368.27	0.08	2012.57	25473.07	0.08	16.42	4.33	79.25	
11	Grain yield per plant (g)	88.95	296.52	0.30	355.08	1186.09	0.30	12.18	18.20	69.62	
12	Shelling percentage	75.93	84.94	0.89	303.73	339.74	0.89	15.43	18.56	66.01	