

# Electronic Journal of Plant Breeding



## Research Article

### Genetic variability analysis for yield and yield attributes among determinate “Wal” type Indian bean [*Lablab purpureus* (L.) sweet] genotypes

D. P. Patel<sup>1\*</sup>, K. G. Modha<sup>1</sup>, A. D. Kyada<sup>2</sup>, Jwala Pranati<sup>3</sup>, M. R. Prajapati<sup>1</sup>, B. H. Kale<sup>1</sup> and R. K. Patel<sup>1</sup>

<sup>1</sup>Department of Genetics and Plant Breeding, N. M. College of Agriculture, Navsari Agricultural University, Navsari-396 450, India.

<sup>2</sup>Division of Genetics, ICAR-Indian Agricultural Research Institute, New Delhi-110012, India.

<sup>3</sup>Department of Genetics and Plant Breeding, Professor Jayashankar Telangana State Agriculture University, Rajendranagar, Hyderabad-500030, India.

\*E-Mail: dhrumilpatel5292@gmail.com

#### Abstract

Fifty two genotypes of Indian bean were examined for the genetic variability parameters and trait correlation analyses for 11 yield contributing traits in  $F_5$  progenies. Analysis revealed that these genotypes have a considerable level of diversity. The magnitude of GCV and PCV was high for seed yield per plant, which indicated the potential for selection based genetic enhancement. The GCV and PCV values were moderate for pod weight, pods width and pod length. High heritability along with high genetic advance as per cent mean was recorded for pod width, pod length, pod weight and seed yield per plant. The seed yield per plant had a significant positive correlation with pod weight, plant height, pod length, pod width and seeds per pod. The overall genotypic path analysis revealed that racemes per plant exhibited the highest positive direct effects on seed yield per plant followed by direct effects of pod length, pods per raceme, pod width and pod weight. Therefore, the simple selection is advised for pod length, pod width and pod weight to boost the yield of the Indian bean. Based on mean values, the progenies W-19-26, W-19-22 and W-19-25 of GNIB-21 × GP-167 among all crosses had shown significantly higher seed yield over the check varieties GNIB-21 and GNIB-22 and hence identified as elite genotypes.

**Keywords:** Indian bean, Variability, Correlation, Heritability, Path analysis

#### INTRODUCTION

Indian bean, [*Lablab purpureus* (L.) sweet] belongs to the Fabaceae family and have varying diploid chromosome number  $2n= 20, 22$  or  $24$ . It was originated in India and introduced from Southeast Asia to Africa in the 8<sup>th</sup> century (Deka and Sarkar, 1990). It is also known as lablab bean, labian bean, Egyptian bean, hyacinth bean, field bean, Dolichos bean, poor man's bean, country bean, bonevise bean, tonga bean and butter bean. In Gujarat and Maharashtra, it is known as a 'Wal'. Throughout the tropics

and subtropics, it is grown as grain and green vegetables as well as fodder. The green and dried pods of Indian beans are enriched with nutrients like carbohydrates, proteins, fat, fibres and minerals viz., Ca, P and Fe (Gopalan *et al.*, 1982 and Naeem *et al.*, 2009). Seeds are rich in protein (15-25 %) as compared to pods (10-19 %) (Gopalan *et al.*, 1982). In Gujarat, many varieties of Indian beans are available for cultivation. Each of these varieties is known for its unique characteristics

like pod shape, size, colour and aroma having a broad genetic base. It is photoperiod sensitive, however, rare photo-insensitive types are also available. In Gujarat, on the basis of pod characteristics, Indian bean cultivars are locally classified as *Wal* (bitter/sweet in-taste, short and flat shape, high fibre content), *Valor* (bitter or sweet in-taste, long and narrow shape, medium fibre content) and *Papdi* (sweet in-taste, short and narrow shape, less fibre content) (Kyada *et al.*, 2022). Owing to its drought tolerance, it can be grown efficiently in a dry land with limited rainfall. Indian bean varieties with determinate growth habits are preferred as they have certain advantages over indeterminate types *viz.*, inter-cropping, early flowering, synchronise maturity and easy and feasible mechanical harvesting

For any genetic improvement programme, the presence of genetic variability is prerequisite in the base population. Highly heritable characters are required for the success of crop improvement programmes. The knowledge on genetic advance and heritability of the desirable traits helps plant breeders to establish accurate and appropriate breeding methodology (Dewangan *et al.*, 2017). The efficiency of selection method is determined by the nature and magnitude of the interaction between heritable and non-heritable variables. Dividing the correlation coefficient into the respective direct and indirect effects would offer

a more relevant explanation on the path with which the independent variables are influencing the dependent variable. Path analysis, a standardized partial regression coefficient, partitions the correlation coefficient into the component of direct and indirect effects and unravels whether the association of these characters with yields is due to their indirect effects via some other trait or traits. The path coefficient explains the cause and effect relation, and also quantifies the relative impact of all the variables (Wright, 1921). Keeping in view the importance of Indian bean and various factors affecting its yield, the present investigation was carried out with an aim to study genetic variability for yield and yield attributing traits in the Indian bean.

## MATERIALS AND METHODS

The present experiment was carried out at College Farm, N. M. College of Agriculture, Navsari Agricultural University, Navsari (Gujarat) during *Rabi*, 2019-20. The experimental material involved fifty-two "*Wal*" type determinate genotypes and out of which, fifty were F<sub>5</sub> progenies and two cultivars GNIB-21 and GNIB-22 were the check varieties. The progenies were obtained from the four crosses *i.e.*, GNIB-21 × GP-1, GNIB-21 × GP-167, GNIB-21 × GP-189, and GNIB-21 × GPKH-120 (**Table 1**) and salient features of parents showed in **Table 2**. All 52 progenies were analyzed in a randomized block design

**Table 1. Progenies of each cross evaluated in the present study**

Cross	Number of progenies	Progenies
GNIB-21 × GP-1	7	W-19-1, W-19-2, W-19-3, W-19-4, W-19-7, W-19-11, W-19-33
GNIB-21 × GP-167	32	W-19-5, W-19-6, W-19-8, W-19-9, W-19-10, W-19-12, W-19-13, W-19-14, W-19-15, W-19-16, W-19-17, W-19-18, W-19-19, W-19-20, W-19-21, W-19-22, W-19-23, W-19-24, W-19-25, W-19-26, W-19-27, W-19-35, W-19-36, W-19-37, W-19-38, W-19-39, W-19-40, W-19-41, W-19-42, W-19-43, W-19-44, W-19-45
GNIB-21 × GP-189	11	W-19-28, W-19-29, W-19-30, W-19-31, W-19-32, W-19-34, W-19-46, W-19-47, W-19-48, W-19-49, W-19-50
GNIB-21 × GPKH-120	0	-

**Table 2. Salient features of parents used in the present study**

S. No.	Parents	Salient features
1.	GNIB-21	Photoperiod insensitive flowering; Determinate and pole type growth habit; No primary branches; Non-pigmented plant; White flowers; Dark green foliage; Prominent grains present inside the pods; Lowest attack of pulse beetle; C shape pods; Creamish white seeds
2.	GP-1	Photoperiod sensitive flowering; Indeterminate growth habit; Light pigmentation on flowers only; Sparsely pigmented stem; Light green, short and broad pods with 3 seeds; Dense foliage; Black seeds
3.	GP-167	Photoperiod sensitive flowering; Indeterminate growth habit; Pigmented buds and flower; Sparsely pigmented stem; Broad leaves; Dark green, long and flat pods, Creamish white bold seeds
4.	GP-189	Photoperiod sensitive flowering; Indeterminate growth habit; Darkly pigmented stem, bud, pod and flower; Purple flowers; Dark green foliage; Pigmented venation, smooth pod surface, Dark purple long narrow pods; Brown seeds with black spots
5.	GPKH-120	Photoperiod sensitive flowering; Indeterminate growth habit; Non pigmented stem, bud, pod, leaves and flower; White flowers; Light green pods with smooth surface; Mosaic susceptible; Dense canopy, Light green long and narrow pods; Creamish white wrinkled seeds

(RBD) with three replications. Each row was comprised of fifteen plants at a spacing of 60 cm and 20 cm between the plants within row. The crop was well grown by providing all basic agronomic and plant protection measures. Total eleven quantitative traits *i.e.*, plant height (PH), days to 50 % flowering (DFF), days to maturity (DM), racemes per plant (RPP), pod weight (PW), pods per plant (PPP), pods per raceme (PPR), seeds per pod (SPP), pod width (PWD), pod length (PL), and seed yield per plant (SYPP) were studied. These abbreviations are used in further discussion for simplicity. The observations were taken from ten randomly selected plants from each progeny and each replication and their means were used in the biometrical analysis. However, days to maturity and days to 50 % flowering were evaluated on a population basis.

Analysis of variance was done as suggested by Panse and Sukhatme (1985). Broad sense heritability ( $h_{bs}^2$ ) and the genetic advance were calculated using

the method indicated by Allard (1999). Genotypic correlation coefficients were calculated using the method recommended by Miller *et al.* (1958). The path analysis was carried out according to Dewey and Lu (1959).

## RESULTS AND DISCUSSION

The analysis of variance for eleven quantitative characters is presented in **Table 3**. This shows significant mean square values for all the characters in  $F_5$  progenies indicating enough variability present among the progenies for all the traits. Similar results were observed by Hadavani *et al.* (2018), Peer *et al.* (2018) and Gamit *et al.* (2020).


Various variability parameters are depicted in **Table 3** and **Graph 1**. The mean values indicated a vast range for DM (84.00-90.33), PH (42.93-70.42 cm), PPP (16.33-49.01), RPP (5.42-11.05), PPR (3.96-7.32), PW (0.73-1.58 g), PL (4.70-8.45 cm), PWD (1.04-2.17 cm) and SYPP (5.25-19.83 g). Based on mean values, the progenies W-19-26

**Table 3. Analysis of variance for eleven quantitative characters under study**

Source of variation	d.f.	Mean sum of square										
		DFF	DM	PH	RPP	PPP	PW	PPR	PL	PWD	SPP	SYPP
Replication	2	0.231	0.231	296.135**	3.599	180.349**	0.379**	5.381**	0.744**	0.014	0.002	8.408
Progenies	51	2.218*	7.274**	118.004**	3.897**	87.031**	0.094**	1.318**	1.469**	0.171**	0.149**	49.214**
Error	102	1.335	0.459	19.851	2.252	37.291	0.014	0.649	0.066	0.005	0.016	730.33

\*\* - Significant at 1.0 % level of probability, \* - Significant at 5.0 % level of probability

DFF=Days to 50% flowering      DM=Days to maturity      PH=Plant height (cm)      RPP=Racemes per plant  
 PPP=Pods per plant      PW=Pod weight (g)      PPR=Pods per raceme      PL=Pod length (cm)  
 PWD=Pod width (cm)      SPP=Seeds per pod      SYPP=Seed yield per plant (g)



**Graph 1. Graphical representation of genetic variability parameters for eleven quantitative characters in  $F_5$  progenies**

GCV (%) = Genotypic coefficient of variation  
 GAM (%) = GA as per cent of mean

PCV (%) = Phenotypic coefficient of variation  
 $h_{bs}^2$  (%) = Heritability (Broad sense)

(19.83 g), W-19-22 (19.45 g) and W-19-25 (19.42 g) of GNIB-21 × GP-167 cross among all crosses had shown significantly higher seed yield over the check varieties GNIB-21 (6.86 g) and GNIB-22 (10.05 g) and hence identified as elite genotypes.

Progenies exhibited higher values for GCV and PCV for SYPP (29.12 % and 35.79 %, respectively). The narrow gap between GCV and PCV indicated less influence of environment on traits *viz.*, DM, PWD, PL and SPP unveiling the scope of improving these characters by selection (Table 4). PPP (13.77 % and 24.82 %, respectively), PW (13.22 % and 16.33 %, respectively), PL (11.24 % and 12.01 %, respectively) and PWD (13.33 % and 13.96 %, respectively) had moderate GCV and PCV values, which indicated that these traits would show less response to selection. Moreover, the magnitude of GCV and PCV values were low for DFF (1.23 % and 2.90 %, respectively), SPP (5.40 % and 6.31 %, respectively) and DM (1.76 % and 1.93 %, respectively) except PH (9.62 % and 12.19 %, respectively). PPR (8.19 % and 16.21 %, respectively) showed moderate PCV values stipulating a narrow range of variability for this trait, which restricts the scope of selection for it. The RPP (0.51 % and 21.48 %, respectively) showed lower GCV and higher PCV values. Similar results are also there in earlier studies for DFF and DM by Asaduzzaman *et al.* (2015); SYPP by Peer *et al.* (2018); PL Choudhary *et al.* (2016); PWD by Noorjahan *et al.* (2019); PW and SPP by Afsan and Roy (2020).

In addition, it indicated moderate to high heritability (broad sense) for all the quantitative traits studied (Table 4). High heritability values for DM (83.17 %), PH (62.24 %), PW (65.52 %), PL (87.57 %), PWD (91.09 %), SPP (73.31 %) and SYPP (66.19 %) indicated that these traits are under little influence of environment, so phenotype could be a good indicator of genotypic effect and traits

could be improved by simple phenotypic selection. The PPP had a moderate heritability (30.78 %) suggesting a medium influence of environment on the expression of this trait and limited scope for phenotypic selection. Low heritability values were depicted by DFF (18.05 %), RPP (19.58 %) and PPR (25.56 %) indicating the major effect of environmental effects on the total phenotypic variation. So, direct selection for these traits is not suggested. Similar results were observed for PH (92.60 %), DM (97.00 %) and SYPP (83.50 %) by Hadavani *et al.* (2018) in 50 genotypes; PWD (85.90 %) and PL (97.50 %) by Noorjahan *et al.* (2019) in 31 genotypes; pods per plant (49.83 %) by Singh *et al.* (2015) in 24 genotypes, and SPP (93.68 %) and PW (95.77 %) by Afsan and Roy (2020) in 11 genotypes.

A high magnitude of genetic advance as per cent of mean (GAM) was found for SYPP (48.80 %), PW (22.03 %), PL (21.67 %) and PWD (26.20 %) (Table 4). Moderate GAM was observed for PH (15.63 %) and PPP (15.73 %), while low estimates of GAM were recorded by DM (3.30 %), DFF (1.08 %), RPP (8.66 %), PPR (8.53 %) and SPP (9.53 %). Similar results were recorded for days to maturity and DFF by Mohan *et al.* (2014); PL, PWD and PW by Noorjahan *et al.* (2019a); SPP by Gupta *et al.* (2010) and SYPP by Peer *et al.* (2018).

Generally, heritability coupled with GAM is more desirable than heritability alone in predicting the response to selection. PW (65.52 % and 22.03 %, respectively), PL (87.57 % and 21.67 %, respectively), PWD (91.09 % and 26.20 %, respectively) and SYPP (66.19 % and 48.80 %, respectively) manifested high heritability coupled with high GAM pointing out the role of additive gene effects. Thus, improvement of these traits could be achieved through direct phenotypic selection. High heritability along with moderate GAM was recorded for PH (62.24 % and 15.63

**Table 4. Genetic variability parameters of quantitative characters studied**

Character	Range		$\sigma^2_E$	$\sigma^2_P$	GCV (%)	PCV (%)	$h^2_{bs}$	GAM (%)
	Minimum	Maximum						
Days to 50 % flowering	42.33	46.00	0.29	1.63	1.23	2.90	18.05	1.08
Days to maturity	84.00	90.33	2.27	2.73	1.76	1.93	83.17	3.30
Plant height (cm)	42.93	70.42	32.72	52.57	9.62	12.19	62.24	15.63
Racemes per plant	5.42	11.05	0.55	2.80	0.51	21.48	19.58	8.66
Pods per plant	16.33	49.01	16.58	53.87	13.77	24.82	30.78	15.73
Pod weight (g)	0.73	1.58	0.03	0.04	13.22	16.33	65.52	22.03
Pods per raceme	3.96	7.32	0.22	0.87	8.19	16.21	25.56	8.53
Pod length (cm)	4.70	8.45	0.47	0.53	11.24	12.01	87.57	21.67
Pod width (cm)	1.04	2.17	0.05	0.06	13.33	13.96	91.09	26.20
Seeds per pod	3.21	4.51	0.04	0.06	5.40	6.31	73.31	9.53
Seed yield per plant (g)	5.25	19.83	14.01	21.18	29.12	35.79	66.19	48.80

GCV (%) = Genotypic coefficient of variation  
 $h^2_{bs}$  (%) = Heritability (Broad sense)

PCV (%) = Phenotypic coefficient of variation  
 GAM (%) = GA as percent of mean

%, respectively) stipulating that this trait is governed by non-additive gene effects and there is a limited scope of selection for improvement. High heritability with low GAM was observed for SPP (73.31 % and 9.53 %, respectively) and DM (83.17 % and 3.30 %, respectively) which is indicative of non-additive genes. So, it is not possible to improve these traits by simple phenotypic selection. Moderate heritability with moderate GAM was observed for PPP (30.78 % and 15.73 %, respectively). Hence, selection would be fairly effective for the improvement of this trait. Low heritability with low GAM was observed for DFF (18.05 % and 1.08 %, respectively), RPP (19.58 % and 8.66 %, respectively) and PPR (25.56 % and 8.53 %, respectively) indicating higher environmental influence, hence little gain is expected through simple selection for these traits. A similar result was observed for heritability and GAM for SYPP (94.20 % and 68.33 %), PWD (95.33 % and 24.67 %), PL (98.90 % and 65.89 %) and PW (99.19 % and 68.17 %) by Peer *et al.* (2018) in 29 genotypes; DM (93.00 % and 3.07 %) by Mohan *et al.* (2014) in 57 genotypes and SPP (85.54 % and 8.77 %) by Gnanesh *et al.* (2006) in 64 genotypes.

The correlation study revealed that SYPP had a significant positive genotypic correlation with PH (0.357\*\*), PW (0.653\*\*), PL (0.274\*), PWD (0.699\*\*) and SPP (0.480\*\*), which indicated that it is possible to improve these traits simultaneously with SYPP through direct phenotypic selection (Table 5). While it had a non-significant positive correlation with RPP (0.086) which showed less significance of this trait for yield improvement. Similar results were observed for PW by Ali *et al.* (2005) and PH and PL by Choudhary *et al.* (2016). PH had a positive non-significant correlation and negative significant correlation with DM (0.187) and DFF (-0.417\*\*), respectively, which suggested that simultaneous selection for more PH and SYPP may result in early flowering via indirect selection. RPP and PWD had a positive significant association with SPP (0.589\*\* and 0.378\*\*, respectively). SPP was

significantly and negatively correlated with PPR (-0.018) (Table 5).

The genotypic path analysis unveiled that the highest positive direct effects on SYPP were imposed by RPP (2.132) followed by PL (0.934), PPR (0.664), PWD (0.610) and PW (0.046) (Table 6). Hence, selection for such characters will be easy in Indian beans. Similar results were also recorded for RPP and PWD by Magalingam *et al.* (2013); PWD and PL by Singh *et al.* (2011) and PPR by Ravinaik *et al.* (2014). The highest negative direct effect on SYPP was recorded by PPP (-2.339) followed by PH (-1.068), SPP (-0.874), DFF (-0.755) and DM (-0.576) suggesting less importance of these traits in selection for higher seed yield. Similar results were also recorded for DFF and SPP by Kamble *et al.* (2015); DM by Singh *et al.* (2015), and PPP and PH by Ravinaik *et al.* (2014). PH imposed the highest positive indirect effect via PPP on SYPP. PPP and SPP exhibited the highest positive indirect effect on SYPP via RPP. PW, PWD and PL exhibited a positive direct effect on SYPP along with high significant and positive association with SYPP. Therefore, these can be regarded as the most important yield attributing characters. In path analysis, a high residual effect of 0.577 was observed which indicates that more yield related traits are necessary to include in future studies to cover all of the available variations.

The overall study indicated that abundant variability is present among the genotypes for all the characters. The progenies W-19-26, W-19-22 and W-19-25 had shown significantly higher seed yield over the check varieties and hence identified as elite genotypes. The magnitude of GCV and PCV was high for seed yield per plant, which indicated the potential for selection based genetic enhancement. High heritability along with high GAM was recorded for pod width, pod length, pod weight and seed yield per plant. The seed yield per plant had a significant positive correlation with pod weight, plant height, pod

**Table 5. Genotypic correlation coefficients of seed yield per plant with other characters in F<sub>5</sub> progenies**

Character	DFF	DM	PH	RPP	PPP	PW	PPR	PL	PWD	SPP	SYPP
DFF	<b>1.000</b>										
DM	-0.189	<b>1.000</b>									
PH	-0.417**	0.187	<b>1.000</b>								
RPP	-0.273	0.205	0.171	<b>1.000</b>							
PPP	0.073	-0.268	-0.603**	0.491**	<b>1.000</b>						
PW	-0.219	-0.189	0.094	-0.189	-0.056	<b>1.000</b>					
PPR	0.481**	-0.425**	-0.845**	-0.054	0.726**	0.058	<b>1.000</b>				
PL	0.167	-0.022	-0.006	-0.235	0.093	0.548**	0.017	<b>1.000</b>			
PWD	-0.376**	0.126	0.441**	-0.042	-0.165	0.620**	-0.314*	0.529**	<b>1.000</b>		
SPP	-0.489**	0.329*	0.301*	0.589**	-0.018	0.254	-0.376**	0.222	0.378**	<b>1.000</b>	
SYPP	-0.288*	-0.162	0.357**	0.086	-0.068	0.653**	-0.210	0.274 *	0.699**	0.480**	<b>1.000</b>

\*\* - Significant at 1.0 % level of probability, \* - Significant at 5.0 % level of probability

**Table 6. Genotypic path coefficient analysis of component characters towards seed yield per plant in F<sub>5</sub> progenies**

Character	DFF	DM	PH	RPP	PPP	PW	PPR	PL	PWD	SPP	Correlation with seed yield per plant
DFF	-0.755	0.109	0.445	-0.581	-0.170	-0.010	0.319	0.156	-0.229	0.428	-0.288 *
DM	0.143	-0.576	-0.200	0.437	0.627	-0.009	-0.283	-0.021	0.008	-0.288	-0.162 NS
PH	0.315	-0.108	-1.068	0.365	1.411	0.004	-0.561	-0.006	0.269	-0.263	0.357 **
RPP	0.206	-0.118	-0.183	2.132	-1.148	-0.009	-0.036	-0.220	-0.025	-0.515	0.086 NS
PPP	-0.055	0.154	0.644	1.046	-2.339	-0.003	0.482	0.087	-0.100	0.015	-0.068 NS
PW	0.165	0.109	-0.100	-0.403	0.131	0.046	0.039	0.512	0.378	-0.222	0.653 *
PPR	-0.363	0.245	0.903	-0.116	-1.699	0.003	0.664	0.016	-0.191	0.328	-0.210 NS
PL	-0.126	0.013	0.006	-0.501	-0.217	0.025	0.011	0.934	0.322	-0.194	0.274 *
PWD	0.284	-0.007	-0.471	-0.086	0.385	0.028	-0.208	0.494	0.610	-0.330	0.699 **
SPP	0.369	-0.190	-0.321	1.255	0.041	0.012	-0.250	0.207	0.230	-0.874	0.480 *

\*\* - Significant at 1.0 % level of probability, \* - significant at 5.0 % level of probability, Residual = 0.577, Bold diagonal figures are the direct effects

length, pod width and seeds per pod. The overall genotypic path analysis revealed that racemes per plant exhibited the highest positive direct effects on seed yield per plant followed by direct effects of pod length, pods per raceme, pod width and pod weight. The proximity of GCV and PCV estimates, high genetic advance as per cent mean, high heritability, positive and significant correlation and the positive direct effect were observed for pod weight, pod width and pod length. So, these traits would be advantageous in a further breeding program for the improvement of seed yield in Indian beans.

## REFERENCES

- Afsan, N. and Roy, A.K. 2020. Genetic variability, heritability and genetic advance of some yield contributing characters in lablab bean (*Lablab purpureus* L. sweet). *Journal of Biosciences*, **28**(1):13-20. [Cross Ref]
- Ali, F., Sikdar, B., Roy, A.K. and Joarder, O.I. 2005. Correlation and genetic variation of twenty different genotypes of lablab bean, *Lablab purpureus* (L.) Sweet. *Bangladesh Journal of Botany*, **34**(2):125-128.
- Allard, R.W. 1999. *Principles of Plant Breeding*. John Wiley and Sons.
- Asaduzzaman, M., Bhuiyan, M.J.H., Hossain, M.A. and Raffi, S.A. 2015. Genetic variability, heritability and genetic advance of yield and yield contributing characters of lablab bean (*Lablab purpureus* L.). *Progressive Agriculture*, **26**(1):22-25. [Cross Ref]
- Choudhary, J., Kushwah, S.S., Singh O.P. and Naruka I.S. 2016. Studies on genetic variability and character association in Indian bean [*Lablab purpureus* (L.) sweet]. *Legume Research*, **39**(3):336-342. [Cross Ref]
- Deka, R.K. and Sarkar, C.R. 1990. Nutrient composition and anti-nutritional factors of *Dolichos hyacinth bean* L. seeds. *Food Chem.*, **38**(4):239-246. [Cross Ref]
- Dewangan, R., Bahadur, V., Choyal, P., Ramesh, Xaxa, S., Vipul, Sachan, S. and Kerketta, A. 2017. Study on genetic variability, heritability and genetic advance in dolichos bean (*Lablab purpureus* L.) genotypes. *International Journal of Current Microbiology and Applied Sciences*, **6**(8):3228-3232. [Cross Ref]
- Dewey, D.R. and Lu, K.H. 1959. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*, **51**(9):515-518. [Cross Ref]
- Gamit, U.C., Jivani, L.L., Ponkia, H.P., Balas, A. and Vadavia, A.T. 2020. Study of genetic variability and heritability in Indian bean (*Lablab purpureus* L.). *Electronic Journal of Plant Breeding*, **11**(1):328-330. [Cross Ref]
- Gnanesh, B.N., Sekhar, M.R., Reddy, K.R. and Eswarareddy, N.P. 2006. Genetic variability, character association and path analysis of pod yield and its component characters in field bean (*Lablab purpureus* L. sweet). *GEOBIOS*, **33**(1):163-168.
- Gopalan, C.V., Ramasastri, B.Y. and Balasubramaniam, S.C. 1982. Nutritive values of Indian food. National Institute of Nutrition, ICMR, Hyderabad. Pp. 75.

- Gupta, K.K., Sharma, T.R. and Tomar, R. 2010. Genetic variability, heritability and genetic advance in Indian bean [*Lablab purpureus* (L.) sweet]. *JNKVV*, **44**(2):159-162.
- Hadavani Janaki, K., Mehta, D.R., Raval, L.J. and Ghetiya, K.P. 2018. Genetic variability parameters in Indian bean (*Lablab purpureus* L.), *International Journal of Pure Applied Bioscience*, **6**(4):164-168. [Cross Ref]
- Kamble, S.S., Devmore, J.P., Sawardekar, S.V., Bhawe, S.G. and Palshetkar, M.G. 2015. Correlation and path coefficient analysis for yield and yield components in segregating ( $F_4$ ) generation of lablab bean (*Lablab purpureus* L. Sweet). *International Journal of Applied Biology and Pharmaceutical Technology*, **6**(3):237-240. [Cross Ref]
- Kyada, A.D., Kale, B.H., Pranati, J., Patel, G.M., Patel, D.P., Prajapati, M.R., Modha, K.G. and Patel, R.K. 2022. Genetic variability, character association and path coefficient analysis in determinate  $F_5$  progenies of Indian bean [*Lablab purpureus* (L.) Sweet]. *Electronic Journal of Plant Breeding*, **13**(2):319-324. [Cross Ref]
- Magalingam, V., Yassin, M. and Kumar, S.R. 2013. Genetic variability and character association in dolichos bean. *SAARC Journal of Agriculture*, **11**(2):161-171. [Cross Ref]
- Miller, P.J., Willams, J.C., Robinsons, H.F. and Comstock, R.E. 1958. Estimates of genotypic and environmental variances and covariance in upland cotton and their implications in selection. *Agronomy Journal*, **50**(3):126-131. [Cross Ref]
- Mohan, N., Aghora, T.S. and Wani, M.A. 2014. Studies on genetic variability in dolichos bean (*Lablab purpureus* L.). *Journal of Horticultural Sciences*, **9**(1):82-85.
- Naeem, M., Khan, M.M.A. and Morris, J.B. 2009. Agrobotanical attributes, nitrogen-fixation, enzyme activities, photosynthesis, nutraceuticals of hyacinth bean (*Lablab purpureus* L.): A bio-functional medicinal legume. *Am J Plant Physiol.*, **4**(2):58-69. [Cross Ref]
- Noorjahan, A.M., Deshmukh, J.D., Wankhade, M.P. and Kalpande H.V. 2019. Genetic variability, heritability and genetic advance studies in dolichos bean (*Lablab purpureus* L.). *International Journal of Chemical Studies*, **7**(3):479-482.
- Panase, V.G. and Sukhatme, P.V. 1985. Statistical methods for agricultural workers. *Indian Council of Agricultural Research*, New Delhi. Pp 134-192.
- Peer, S.S., Reddy, P.S.S., Sadarunnisa, S., Reddy, D.S. and Pandravada, S.R. 2018. Genetic variability and heritability for yield and yield attributes in field bean (*Lablab purpureus* L.) genotypes. *International Journal of Current Microbiology and Applied Sciences*, **7**(3):2131-2137. [Cross Ref]
- Ravinaik, K., Hanchinamani, C.N., Patil, M.G. and Imamsaheb, S.J. 2014. Correlation and path coefficient analysis in dolichos bean (*Dolichos lablab* L.) genotypes. *The Asian Journal of Horticulture*, **9**(2):396-399. [Cross Ref]
- Singh, P.K., Rai, N., Lal, H., Bhardwaj, D.R., Singh, R. and Singh, A. 2011. Correlation, path and cluster analysis in hyacinth bean (*Lablab purpureus* L. sweet). *Journal of agricultural Technology*, **7**(4):1117-1124.
- Singh, S., Singh, P.K., Singh, D.R., Pandey, V.B. and Srivastava, R.C. 2015. Genetic variability and character association study in dolichos bean. *Indian Journal of Horticulture*, **72**(3):343-346. [Cross Ref]
- Wright, S. 1921. Correlation and causation. *Journal of Agricultural Research*, **20**:557-587.