

Electronic Journal of Plant Breeding



Research Note

Marker assisted selection and genome recovery for heat tolerance in segregants of rice (*Oryza sativa* L.)

V. Silpa¹, P. Sindhumole^{1&2*}, Jiji Joseph¹, Deepu Mathew³ and Rose Mary Francies⁴

¹Department of Plant Breeding & Genetics, College of Agriculture, Vellanikkara, Kerala Agricultural University, Thrissur-680 656, Kerala, India.

²AICRP on Medicinal & Aromatic Plants and Betel vine, College of Agriculture, Vellanikkara, Kerala Agricultural University, Thrissur-680 656, Kerala, India.

³Department of Plant Biotechnology, CPBMB, College of Agriculture, Vellanikkara, Kerala Agricultural University, Thrissur-680 656, Kerala, India.

⁴Agricultural Research Station, Mannuthy, KAU, Thrissur, Kerala, India.

*E-Mail: sindhumole.p@kau.in

Abstract

In the present global warming scenario, it is urgent to impart heat tolerance into the popular high yielding rice (*Oryza sativa* L.) varieties to reduce yield loss to a great extent. Uma (a prevalent red-grained rice variety in Kerala) was crossed with Nagina 22 (N22), the popular donor for heat tolerance. From the seven promising heat-tolerant F₃ plants previously identified through marker assisted selection and field screening, F₄ lines (58 nos.) were raised and characterised morphologically. Heat tolerance was scored under natural heat stress during hot summer months in field conditions. Among the evaluated F₄ progenies, 26 promising plants registered >75% spikelet fertility. Double panicles were observed in some of the tillers of fourteen F₄ plants. During the selection with reported polymorphic markers between parents, all the 26 F₄ plants produced the corresponding amplicon for RM5749 (linked marker for heat tolerance in the parental cross Uma x N22) and also for RM9, RM201, RM208, RM225, RM242, RM495, RM3586, RM6100, RM6836, and RM26212, corresponding to the heat tolerant parent N22. Genomic analysis revealed 64% recovery of the Uma genome in seven F₄ plants, with maximum genomic regions of Uma on the chromosomes 3 and 5. These promising F₄ plants could be forwarded to generate stable and high yielding heat tolerant rice varieties. The identified markers could be used for linkage analysis and QTL mapping in future.

Keywords: Heat stress; Rice; spikelet fertility; SSR; QTL

Rice (*Oryza sativa* L.) is the staple food all around the globe, feeding more than one-third of the human population (Sarangi *et al.*, 2019). The global demand for rice will be about 584 million tons towards 2050 (Samal *et al.*, 2022). Climate change is a threat for food security by intensifying the extreme weather events (Venkateshwarlu *et al.*, 2022). Heat stress due to rapid global warming is a serious threat to the world's food security (Xu *et al.*, 2021) and a major challenge to the rice breeders. Though rice

can grow normally at a temperature range of 27 to 32°C, temperature >32°C negatively affects all of its growth and development stages (Aghamolki *et al.*, 2014). The growth, yield, and quality of rice are influenced by high temperature stress, which disrupts its phenology, physiology, and yield components (Sindhumole *et al.*, 2021). In rice, flowering and seed development stages are very sensitive to heat stress, leading to unfilled and chalky grains, which reduce the quantity and quality of seeds.

Genetically improving the crops for heat tolerance is one of the best approaches to tackle this problem (Chidambaranathan *et al.*, 2021). Heat tolerance is a multi-genic trait of which candidate genes are not well known. So, using yield as the selection criterion for developing heat tolerant varieties is usually unreliable during the early stages. Hence, marker assisted selection (MAS) is a better alternative to early generation selection. Fixing the recipient alleles in

non-target regions and identifying the genotypes containing cross-overs close to the target genes are the major advantages of using DNA markers (Tanksley *et al.*, 1989). As several QTLs associated with heat tolerance traits are with complex inheritance, MAS is more efficient for their screening. SSR markers are used for identifying various abiotic stress tolerance traits in rice (Salam *et al.*, 2017; Waghmare *et al.*, 2018).

Table 1. Details of SSR primers used for polymorphism study

S.No.	Primer	Chromo some no.	Sequence		Annealing temperature (°C)	Product size (bp)
			Forward (5'-3')	Reverse (5'-3')		
1	RM212	1	CCACTTTCAGCTACTACCAG	CACCCATTGTCTCTCATTATG	54.5	136
2	RM302	1	TCATGTCATCTACCATCACAC	ATGGAGAAGATGGAATACTTGC	59	156
3	RM495	1	AATCCAAGGTGCAGAGATGG	CAACGATGACGAACACAACC	59	159
4	RM10346	1	GCTTGATCTGCCCTTGTCTTGG	AACTCGAGCGGCCTTCTCAGC	65	292
5	RM9	1	GGTGCCATTGTCTGTCCTC	ACGGCCCTCATCACCTTC	59	136
6	RM3340	2	TCTTGGCAAGCTCTCCTCTC	CCATCATCTCGATCTTGACG	59	117
7	RM166	2	GGTCTGGGTCAATAATTGGGTTACC	TTGCTGCATGATCCTAAACGG	59	321
8	RM208	2	TCTGCAAGCCTGTCTGATG	TAAGTCGATCATTGTGTGGACC	59	173
9	RM251	3	GAATGGCAATGGCGCTAG	ATGCGGTTCAAGATTCGATC	59	147
10	RM85	3	CCAAAGATGAAACCTGGATTG	GCACAAGGTGAGCAGTCC	59	107
11	RM3586	3	GAAGAGAGAGCCAGAGCCAG	ACACGATCGAGCTAGAAGACG	59	118
12	RM280	4	ACACGATCCACTTTGCGC	TGTGTCTTGAGCAGCCAGG	60	155
13	RM252	4	TTCGCTGACGTGATAGGTTG	ATGACTTGATCCCGAGAACG	59	216
14	RM5749	4	GTGACCACATCTATATCGCTCG	ATGGCAAGGTTGGATCAGTC	58	162
15	RM518	4	CTTTCACTCACTCACCATGG	ATCCATCTGGAGCAAGCAAC	59	171
16	RM169	5	TGGCTGGCTCCGTGGGTAGCTG	TCCCGTTGCCGTTTCATCCCTC	71	167
17	RM163	5	ATCCATGTGCGCCTTTATGAGGA	CGTACCTCCTTCACTTACTAGT	59	124
18	RM164	5	TCTTGCCCGTCACTGCAGATATCC	GCAGCCCTAATGCTACAATTCTTC	60	246
19	RM13	5	TCCAACATGGCAAGAGAGAG	GGTGGCATTGATTCCAG	59	141
20	RM6836	6	TGTTGCATATGGTGCTATTTGA	GATACGGCTTCTAGGCCAAA	59.5	240
21	RM225	6	TGCCCATATGGTCTGGATG	GAAAGTGGATCAGGAAGGC	58.5	140
22	RM755	6	AAAGGATAAATGTGGGGATC	ATAACCGTCTGGTTTCACTG	54.5	141
23	RM336	7	CTTACAGAGAAACGGC ATCG	GCTGGTTTGTTCAGGTTTCG	58	154
24	RM447	8	CCCTTGTGCTGTCTCCTCTC	ACGGGCTTCTTCTCCTTCTC	58	111
25	RM256	8	GACAGGGAGTGATTGAAGGC	GTTGATTTGCCAAGGGC	58	127
26	RM242	9	GGCCAACGTGTGTATGTCTC	TATATGCCAAGACGGATGGG	52.5	225
27	RM201	9	CTCGTTTATTACCTACAGTACC	CTACCTCCTTTCTAGACCGATA	54.5	158
28	RM6100	10	TCCTCTACCAGTACCGCACC	GCTGGATCACAGATCATTGC	58	144
29	RM254	11	AGCCCCGAATAAATCCACCT	CTGGAGGAGCATTGGTAGC	60	165
30	RM552	11	CGCAGTTGTGGATTTCAGTG	TGCTCAACGTTTGACTGTCC	58	195
31	RM26212	11	GTCGCTCCTCTCCTCCAATCC	GCTCGCTGCTCTAATCTCTTGC	62.5	180
32	RM3701	11	GAGCTAGAGGGAGGAGGTGC	TTGACTGATAGCCGATTGGG	62.5	174
33	RM224	11	ATCGATCGATCTTCACGAGG	TGCTATAAAAGGCATTCCGGG	59	157
34	RM19	12	CAAAAACAGAGCAGATGAC	CTCAAGATGGACCCAAGA	58	226
35	RM17	12	TGCCCTGTTATTTCTTCTCTC	GGTGATCCTTTCCATTT CA	59	184

Resorting to both genotypic and phenotypic selection gives a better output in breeding programmes. In Kerala, where rice is a major crop, temperature $>39^{\circ}\text{C}$ causes heat stress-induced spikelet sterility. Among the prominent varieties of the state, Uma is high yielding but very susceptible to heat during reproductive and grain filling phases. To enhance heat tolerance in rice, the rice variety Nagina 22 (N22), which is characterized by better panicle length, panicle weight, and 1000-grain weight under heat stress variety is being used as a donor in several breeding programmes (Vinitha *et al.*, 2020).

To incorporate high temperature tolerance, Uma rice variety was crossed with N22 (donor for heat tolerance), and the segregating generations were subjected to early MAS to identify superior progenies with heat tolerance. The present study was undertaken to evaluate the morphological variations in the F_4 population of Uma x N22, and to analyse the allelic distribution of various polymorphic marker loci identified between the parents. F_4 progenies (58 nos.) raised from the selected seven heat tolerant F_3 plants of the cross Uma x N22 were evaluated in pots inside the poly house in the field lab of Department of Plant Breeding and Genetics, College of Agriculture, Vellanikkara, KAU, Thrissur. This unreplicated experiment was conducted during the summer months of 2018. Hence, the plants were stressed with high temperature ($>40^{\circ}\text{C}$), especially during the reproductive phase, and the biometrical observations were recorded.

Genotyping: Genomic DNA of F_4 segregants and the parents were extracted as per the CTAB method (Murray and Thompson, 1980). DNA purity was checked by measuring the OD values at 260 and 280 nm using NanoDrop ND100 spectrophotometer (Jenway-

Genova Nano). For studying the allelic distribution in F_4 generation, 35 SSR markers polymorphic between Uma and N22 (Waghmare *et al.*, 2018) were used (**Table 1**). Polymerase Chain Reaction (PCR) was performed with markers in a thermal cycler (Eppendorf Mastercycler® nexus gradient PCR), and the amplified products were subjected to electrophoresis in 2% Agarose gel.

Morphological evaluation: Biometric characters of 58 F_4 segregants were recorded and furnished in **Table 2**. Spikelet fertility is the major trait for the identification and selection of plants for heat tolerance. In the current study, heat tolerant parent N22 exhibited the highest spikelet fertility (90.13%), while the heat sensitive parent Uma exhibited a lower spikelet fertility (55.29%) during summer months. Twenty six F_4 plants, showing more than 75% spikelet fertility in the field, were selected (**Table 3** and **Fig. 1**). Time taken for the appearance of first panicle in selected F_4 plants was equivalent to that of N22 (76 days), whereas no. of filled grains per panicle (79.12) was lesser than both the parents. Mean grain weight per plant of the selected F_4 plants was 5.82 g, comparable to Uma.

Usually, only a single panicle is produced in each productive tiller of rice. In contrast, double panicles were observed in some tillers of 14 F_4 segregants in this study (**Table 2**). This phenomenon was observed in single tiller of nine plants (Plant Nos. 15.3, 15.4, 16.5, 16.6, 16.9, 31.4, 31.5, 41.1 and 41.2), in two tillers of four plants (Plant Nos. 13.1, 16.3, 16.4 and 31.10) and in three tillers of one plant (Plant No. 31.3). Double panicles per productive tiller was also observed earlier by Sindhumole (unpublished) in the parental F_2 (Uma x N22) plant, from which F_4 plants of the present study were derived. Similarly, Anup (2022) observed double panicles per tiller in N22 variety while




Fig. 1. Variability of spikelet fertility (%) in F_4 plants

Table 2. Morphological characters of F₄ segregants of Uma x N22 rice hybrid

S.No.	Plant no.	Tillers/ plant	Panicles/ plant	Panicle length (cm)	Days for flowering	Spikelets/ panicle	Filled grains/ panicle	Partially filled grains / panicle	Sterile grains/ panicle	100 seed wt (g)	Grain wt/ plant (g)	Spikelet fertility (%)
1	12.1	10	8	19.5	70	121.1	12.2	32.6	76.3	1.50	3.10	37.00
2	12.2	14	5	16.0	69	111.8	23.8	35.4	52.6	2.50	4.30	52.96
3	12.3	10	8	18.5	70	89.2	15.2	36.4	37.6	1.90	3.80	57.85
4	12.4	11	11	19.0	75	128.2	31.0	50.2	47.0	2.00	4.60	63.34
5	13.1	8	10 [#]	20.5	69	149.4	32.4	54.6	62.4	2.30	5.90	58.24
6	13.2	12	7	19.2	68	121.7	25.8	35.4	60.5	2.70	5.30	50.29
7	13.3	10	8	17.0	69	109.0	17.4	58.2	33.4	1.90	4.80	69.36
8	15.1	11	10	22.5	85	138.0	97.3	12.4	28.3	1.94	7.30	79.50
9	15.2	10	8	19.6	82	111.2	90.6	8.2	12.4	1.99	7.10	88.85
10	15.3	8	9 [#]	20.5	80	164.8	108.6	27.8	28.4	1.85	7.60	82.77
11	15.4	9	10 [#]	21.2	81	109.0	83.2	10.4	15.4	2.01	7.08	85.88
12	15.5	10	8	21.0	81	129.3	75.4	21.2	32.7	1.58	4.01	74.71
13	15.6	9	9	19.8	80	69.2	30.0	13.8	25.4	2.12	5.80	63.30
14	15.7	9	8	22.0	80	97.8	53.6	26.0	18.2	1.73	6.60	81.40
15	15.8	10	10	21.4	79	37.8	12.0	4.0	21.8	1.91	6.90	42.33
16	16.1	7	7	20.2	69	127.0	105.4	12.0	9.6	1.98	8.21	92.45
17	16.2	8	8	19.1	73	118.7	84.0	20.5	14.2	1.94	7.30	88.04
18	16.3	9	11 [#]	20.5	75	129.6	100	6.0	23.6	2.11	7.50	81.80
19	16.4	8	10 [#]	20.2	70	117.6	73.2	29.4	15.0	1.68	6.51	87.25
20	16.5	8	9 [#]	22	73	117.5	62.8	38.0	17.5	1.45	4.53	85.11
21	16.6	7	8 [#]	18.6	75	130.6	82.0	16.4	32.2	1.90	3.16	75.35
22	16.7	9	9	19.0	76	118.4	78.0	26.4	14.0	2.70	4.64	88.18
23	16.8	8	6	22.0	76	175.2	80.0	22.0	73.2	2.22	5.93	58.22
24	16.9	7	8 [#]	19.5	75	179.3	88.3	21.0	70.0	1.67	4.50	60.96
25	16.10	8	8	18.8	75	147.5	85.4	23.0	39.1	1.75	4.11	73.50
26	31.1	8	7	19.5	81	105.3	48.6	34.2	22.5	2.04	6.10	78.64
27	31.2	10	8	19.6	77	136.6	108.0	18.4	10.2	2.35	7.91	92.54
28	31.3	8	11 [#]	20.0	78	149.4	104.4	31.2	13.8	1.97	5.70	90.77
29	31.4	9	10 [#]	23.0	77	139.0	91.0	32.6	15.4	1.90	6.81	88.93
30	31.5	8	9 [#]	19.5	75	132.8	80.2	41.6	11.0	1.95	6.54	91.72
31	31.6	10	9	19.2	79	174.4	130.2	8.0	36.2	1.94	8.10	79.25
32	31.7	9	8	19.7	77	42.2	28.0	5.6	8.6	2.12	5.80	79.63
33	31.8	9	7	22.0	80	106.8	63.2	26.0	17.6	1.70	4.57	83.53
34	31.9	8	7	23.5	75	124.8	79.4	15.4	30.0	1.30	5.91	75.97
35	31.10	7	9 [#]	19.5	79	114.6	59.0	34.8	20.8	2.60	4.98	81.85
36	31.11	8	8	19.8	77	93.9	61.5	13.0	19.4	2.40	6.53	79.34
37	31.12	12	10	17.0	83	128.8	90.6	23.8	14.4	1.8	7.30	88.82
38	41.1	7	8 [#]	20.8	81	141.0	0.0	28.0	113.0	0.00	0.00	19.86
39	41.2	8	9 [#]	19.2	81	161.0	16.6	8.2	136.2	1.79	3.65	15.41
40	41.3	7	6	22.0	78	166.2	0.0	0.0	166.2	0.00	0.00	0.00
41	41.4	8	8	20.9	81	192.4	79.2	31.8	81.4	1.84	4.90	57.70
42	41.5	8	7	23.7	80	133.8	75.8	34.5	23.5	1.64	4.70	82.44
43	41.6	9	8	22.4	76	149.6	18.6	110.6	20.4	1.32	3.95	86.37
44	41.7	8	7	18.6	76	121.8	12.6	32.8	76.4	2.41	4.55	37.28

S.No.	Plant no.	Tillers/ plant	Panicles/ plant	Panicle length (cm)	Days for flowering	Spikelets/ panicle	Filled grains/ panicle	Partially filled grains / panicle	Sterile grains/ panicle	100 seed wt (g)	Grain wt/ plant (g)	Spikelet fertility (%)
45	41.8	8	8	22.4	83	109.1	17.5	58.0	33.6	1.84	4.29	69.21
46	41.9	9	8	17.0	85	92.9	15.0	47.2	30.7	1.36	4.07	66.96
47	41.10	7	8	20.6	80	146.1	28.0	71.6	46.5	2.75	5.01	68.18
48	41.11	8	8	22.6	79	165.6	45.0	30.2	90.4	3.71	5.40	45.42
49	41.12	10	7	18.8	79	149.3	32.4	54.0	62.9	1.96	4.28	57.81
50	45.1	8	8	19.6	77	139.4	23.2	5.4	110.8	1.80	3.00	20.52
51	45.2	8	7	19.8	79	144.3	15.6	8.5	120.2	1.10	3.10	16.71
52	45.3	9	8	20.5	77	231.0	99.2	20.4	111.4	1.70	3.80	51.78
53	45.4	9	8	17.9	76	102.0	8.0	13.1	80.9	0.80	3.50	20.69
54	45.5	8	7	22.2	76	138.0	9.0	36.4	92.6	0.50	1.20	32.90
55	45.6	11	8	18.5	79	72.0	14.8	21.0	36.2	2.20	3.90	49.73
56	45.7	9	8	16.0	80	90.6	44.2	16.4	30.0	1.10	4.20	66.89
57	45.8	10	8	20.5	80	192.4	79.2	31.8	81.4	1.90	5.80	57.70
58	45.9	9	8	18.4	79	83.6	40.2	15.0	28.4	1.40	3.60	66.03

*Plant with double panicles in some of its tillers

evaluating N22, Jyothi and their hybrid. Hence, this trait of double panicles per tiller in segregants of Uma x N22 might have transferred from N22. However, exact reason for the appearance of double panicles per tiller in rice plants could not be traced from the available literature.

Among the selected 26 F₄ plants (with >75% spikelet fertility), ten (15.3, 15.4, 16.3, 16.4, 16.5, 16.6, 31.3, 31.4, 31.5 and 31.10) had double panicles, but others were with single panicle per tiller. Moreover, though other four plants (13.1, 16.9, 41.1 and 41.2) had double panicles, their spikelet fertility was not up to the desired level (Table 2).

Genotyping: In recent years, genotyping using SSR markers has been proven as the valuable tool for uncovering the hidden information in plants, like domestication features and crop improvement aspects via marker assisted breeding (Badri *et al.*, 2022). Being co-dominant, SSR markers are ideal for segregation analysis in large population sets (Gupta and Varshney, 2000; Guo *et al.*, 2015). Molecular markers are highly effective in assessing genetic diversity, structural analysis, and phylogenetic relations among rice accessions (Bollinedi *et al.*, 2020; Shivani *et al.*, 2021). In marker assisted breeding programmes, a robust marker-trait association should always be preferred, for exploiting a particular marker to improve a specific trait (Bresseghele and Sorrells, 2006).

Foreground selection: Among the segregants of Uma x N22 cross, four F₄ plants (Plant nos. 16.1, 31.2, 31.3, and 31.5) with >90% spikelet fertility and 22 F₄ plants with >75-90% spikelet fertility were subjected to foreground selection along with the parents (N22 and Uma) using

the linked marker RM5749. In parents, the corresponding amplicons (157 bp and 129 bp, respectively) were observed for RM5749, located on chromosome 4 (Fig. 2). Interestingly, all the 26 selected plants registered only one amplicon (157 bp) corresponding to the heat tolerant parent N22, proving that the heat tolerance in these selected lines was inherited from the donor parent N22.

Thirty five SSR markers identified as polymorphic between the two parental lines by Waghmare *et al.* (2018) (Table 1) were used for screening the selected F₄ plants. Fourteen among these 35 SSR markers were recognised as associated with QTL for heat tolerance by earlier researchers (Zhang *et al.*, 2009; Buu *et al.*, 2014 *etc.*). In this study, single marker analysis revealed significant (P<0.05) association of spikelet fertility with RM5749, RM9, RM208, RM201, RM225, RM242, RM495, RM3586, RM6836, RM26212, and RM6100. Poli *et al.* (2013) and Bharathkumar *et al.* (2014) suggested RM225 and RM6100 respectively as polymorphic markers for heat tolerance. RM242 (Wei *et al.*, 2013) and RM3586 (Lang *et al.*, 2015) also were reported as polymorphic markers for heat tolerance.

Xiao *et al.* (2011) reported RM6100 as linked with a major QTL on chromosome 10 for heat tolerance during flowering stage. In conformity with the previous findings of QTL for heat tolerance, the above mentioned eleven markers exhibited the same allelic pattern of the male parent N22 in these 26 fertile F₄ plants studied, indicating the similarity in alleles with heat tolerant parent N22 at these marker loci. These 26 lines were further screened based on genome recovery and trait similarities with parents.




Fig. 2. Foreground selection of 26 fertile F₄ plants using RM5749

RM5749 (chromosome 4) has been identified by Waghmare *et al.* (2018) as co-segregating with heat tolerance in Uma x N22 and its segregants. This emphasizes its efficiency in MAS programs, especially with the segregating generations of the same cross. The position of RM5749 was reported to be at 22.80 Mb on chromosome 4 (Gramene Database). This study also confirms the findings of Buu *et al.* (2014) that RM5749 was linked to heat tolerance in BC₁F₁ lines of the cross OM5930 x N22.

Allele distribution in selected segregants: The present investigation revealed more allelic diversity in the studied segregants (**Table 4**). Among the 34 polymorphic SSR markers used, nine markers (RM13, RM163, RM164, RM166, RM169, RM212, RM302, RM518, and RM7076) exhibited the same allelic pattern of female parent (Uma) in the selected 26 F₄ plants (**Fig. 3**).

Presence of alleles of both the parents was revealed by 17 SSR markers, indicating the heterozygous nature of these marker loci in the selected F₄ segregants, which were

further screened based on genome recovery using the marker data and trait similarities with Uma. Among these, the highest genome recovery of Uma was in plant nos. 16.7(67.14%) and 31.9 (65.71%). Seven plants (Plant no.15.3, 31.3, 31.4, 31.5, 31.7, 31.10, and 41.5) showed 64.29% recovery with Uma, and two plants, 15.4 and 15.7 showed 52.86% and 54.29% recovery of N22 respectively. The GGT(graphical genotyping) software output indicated the maximum genomic regions of Uma on chromosome 3 and chromosome 5 while heterozygous loci were located on chromosomes 3, 5, 11, and 12 (**Fig. 4**).

Thus, during this research programme, the selected F₄ segregants inherited the heat tolerance trait from the donor parent N22. Seven F₄ segregants were identified with 64.29% recovery of the Uma genome (recurrent parent) and the heat tolerance allele related to the donor parent N22. Hence, these promising F₄ lines with better heat tolerance and grain yield can be forwarded to subsequent generations and subjected to phenotyping and genotyping to develop the promising heat tolerant varieties in the background of Uma.

Table 3. Phenotyping of F₄ segregants with more than 75% spikelet fertility

Trait	Maximum	Minimum	Mean
Plant height (cm)	99.0	73.6	81.11
Tillers/plant	12.0	7.0	8.69
Panicles/plant	11.0	7.0	8.65
Panicle length (cm)	23.7	17.0	20.53
Days for flowering	85.0	69.0	77.42
Spikelets /panicle	174.4	42.2	123.54
Filled grains/panicle	130.2	18.6	79.12
Sterile grains/panicle	36.2	8.6	18.94
100 seed weight (g)	2.7	1.3	1.89
Grain weight /plant (g)	7.6	3.1	5.82


Table 4. Distribution of alleles of marker loci used for background selection in the selected F₄ plants

S. No.	RM Marker	N22	15.1	15.2	15.3	15.4	15.7	16.1	16.2	16.3	16.4	16.5	16.6	16.7	31.1	31.2	31.3	31.4	31.5	31.6	31.7	31.8	31.9	31.10	31.11	31.12	41.5	41.6	Uma	
1	RM302	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
2	RM212	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
3	RM495	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
4	RM10346	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
5	RM9	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
6	RM3340	A	A	B	A	H	A	B	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
7	RM166	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
8	RM251	A	A	A	H	A	H	A	A	H	A	A	A	H	H	B	H	H	H	H	H	H	H	H	H	H	H	H	H	B
9	RM208	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
10	RM85	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
11	RM7076	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
12	RM3586	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
13	RM280	A	A	H	B	A	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	B
14	RM252	A	B	A	B	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	B
15	RM518	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
16	RM169	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
17	RM163	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
18	RM164	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
19	RM13	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
20	RM6836	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
21	RM225	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
22	RM7555	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
23	RM366	A	A	A	A	B	A	H	B	H	A	B	H	A	B	H	A	A	A	A	A	A	A	A	A	A	A	A	A	B
24	RM447	A	A	A	B	B	A	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
25	RM256	A	A	B	B	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
26	RM242	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
27	RM201	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
28	RM6100	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
29	RM254	A	B	A	B	A	A	H	B	H	H	H	H	H	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
30	RM552	A	B	A	B	A	A	A	A	A	A	A	A	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
31	RM26212	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B
32	RM3701	A	H	B	A	A	A	H	A	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	B
33	RM224	A	B	B	B	A	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
34	RM19	A	B	B	B	H	H	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
35	RM17	A	H	B	B	A	A	B	B	A	A	A	A	A	H	H	H	H	H	H	H	H	H	H	H	H	H	H	H	B

A – Alleles similar to male parent N22

B – Alleles similar to female parent Uma

H – Heterozygous loci



B: Blank N: Nagina22 U: Uma

Fig. 3. Background selection of 26 fertile F₄ plants using RM518




Fig. 4. GGT software output for the genome of 26 fertile F₄ plants of Uma x N22

REFERENCES

Aghamolki, M. T. K., Yusop, M. K., Oad, F. C., Zakikhani, H., Jaafar, H. Z. and Kharidah, S. 2014. Heat stress effects on yield parameters of selected rice cultivars at reproductive growth stages. *Journal of Food, Agriculture and Environment*, **12**:741-746.

Anup, V. S. 2022. Incorporation of heat tolerance into Jyothi rice (*Oryza sativa* L.) variety through marker assisted breeding. M.Sc. (Ag.) thesis, Kerala Agricultural University, India.

Badri, J., Lakshmidevi G., JaiVidhya, L. R. K., Prasad, M. S., Laha, G. S., Lakshmi, V. J., Isetty, S.R.,

- Padmashree, R., Balakrishnan, D., Varanasi, Y. V. P., Jukanti, A. K., Singh, U. M., Singh V.K., Kumar A., Ram, T., Rao, L. V. S. and Sundaram, R. M. 2022. Multiparent-derived, marker-assisted introgression lines of the elite Indian rice cultivar, 'Krishna Hamsa' show resistance against bacterial blight and blast and tolerance to drought. *Plants*, **11**:622. [Cross Ref]
- Bharathkumar, S., Pragnya, P. J., Jitendra, K., Archana, B., Singh, O. N. and Reddy, J. N. 2014. Identification of rice germplasm associated with microsatellite (SSR) markers for heat tolerance at reproductive stage and expression of heat stress related gene. *Indian Research Journal of Genetics and Biotechnology*, **6**:424-427.
- Bollinedi, H., Yadav, A. K., Vinod, K. K., Krishnan, S. G., Bhowmick, P. K., Nagarajan, M., Neeraja, C. N., Ellur, R. K. and Singh, A. K. 2020. Genome-wide association study reveals novel marker trait associations governing the localization of Fe and Zn in the rice grain. *Frontiers in Genetics*, **11**:213. [Cross Ref]
- Breseghele, F. and Sorrells, M. E. 2006. Association analysis as a strategy for improvement of quantitative traits in plants. *Crop Science*, **46**(3):1323-1330. [Cross Ref]
- Buu, B. C., Ha, P. T. T., Tam, B. P., Nhien, T. T., Hieu, N. V., Phuoc, N. T., Minh, L. T., Giang, L. H. and Lang, N. T. 2014. Quantitative trait loci associated with heat tolerance in rice (*Oryza sativa* L.). *Plant Breeding and Biotechnology*, **2**:14-24. [Cross Ref]
- Chidambaranathan, P., Balasubramaniasai, C., Behura, N., Purty, M., Samantaray, S., Subudhi H., Ngangkham, U., Devanna, B. N., Katara, J. L. and Behera, L. 2021. Effects of high temperature on spikelet sterility in rice (*Oryza sativa* L.): Association between molecular markers and allelic phenotypic effect in field condition. *Genetic Resources and Crop Evolution*, **68**:1923-1935. [Cross Ref]
- Guo, Y., Wu, Y., Anderson, J.A., Moss, J. Q. and Zhu, L. 2015. Disomic inheritance and segregation distortion of SSR markers in two populations of *Cynodon dactylon* (L.) Pers. var. *dactylon*. *PLoS One*, **10**(8):0136332. [Cross Ref]
- Gupta, P. K. and Varshney, R. 2000. The development and use of microsatellite markers for genetic analysis and plant breeding with emphasis on bread wheat. *Euphytica*, **113**(3):163-185. [Cross Ref]
- Lang, N. T., Ha, P. T., Tru, P. C., Toan, T. B., Buu, B. C. and Cho, Y. C. 2015. Breeding for heat tolerance rice based on marker-assisted backcrossing in Vietnam. *Plant Breeding and Biotechnology*, **3**:274-281. [Cross Ref]
- Murray, H. G. and Thompson, W. F. 1980. Rapid isolation of high molecular weight DNA. *Nucleic Acid Research*, **8**:4321-4325. [Cross Ref]
- Poli, Y., Basava, R. K., Panigrahy, M., Vinukonda, V. P., Dokula, N. R., Voleti, S. R., Desiraju, S., and Neelamraju, S. 2013. Characterization of a Nagina22 rice mutant for heat tolerance and mapping of yield traits. *Rice*, **6**(36): 1-9. [Cross Ref]
- Salam, S. A., Sindhumole, P., Waghmare, S. G. and Sajini, S. 2017. Molecular characterization of rice (*Oryza sativa* L.) genotypes for drought tolerance using two SSR markers. *Electronic Journal of Plant Breeding*, **8**(2): 474-479. [Cross Ref]
- Samal, P., Babu, S.C., Mondal, B and Mishra, S. N. 2022. The global rice agriculture towards 2050: An inter-continental perspective. *Outlook on Agriculture*, **51**(2): 164-172. [Cross Ref]
- Sarangi, S., Mandal, C., Dutta, S., Mukherjee, P., Mandal, R., Kumar, S. P. J., Choudhury, P. R., Singh, V. P., Tripathi, D. K. and Mandal, A. B. 2019. Microprojectile based particle bombardment in development of transgenic indica rice involving AmSOD gene to impart tolerance to salinity. *Plant Gene*, **19**:100183. [Cross Ref]
- Shivani, D., Zabeen, F., Chaithanya, K., Koushik, M. B. V. N., Dileep, G. D., Koti, E. P., Supriya K, Sundaram, R. M., Kumar, J. A. and Fiyaz, R. A. 2021. Assessment of genetic diversity of rice germplasm using microsatellite markers. *Pharma Innovation Journal*, **10**(5):1393-1397.
- Sindhumole, P., Silpa, V. and Waghmare, S. G. 2021. Marker assisted breeding for heat tolerance in rice (*Oryza sativa* L.). *Book of Abstracts, International Conference on FCPPB*, 6-7, October, 2021, TNAU, Coimbatore, pp.103.
- Tanksley, S. D., Young, N. D., Paterson, A. H. and Bonierbale, M. W. 1989. RFLP mapping in plant breeding: new tools for an old science. *Biotechnology*, **7**(3):257. [Cross Ref]
- Venkateshwarlu, C., Kole, P. C., Paul, P. J., Singh, A. K., Singh, V. K. and Kumar, A. 2022. Capturing genetic variability and identification of promising drought-tolerant lines in exotic landrace derived population under reproductive drought stress in rice. *Frontiers in Plant Science*, **13**:1-12. [Cross Ref]
- Vinitha, A., Raveendran, M., Prasad, V. B. R. and Vijayalakshmi, D. 2020. Rice yield under high temperature stress is influenced by morphophysiological traits. *Madras Agricultural Journal* **107**(7-9):1-5. [Cross Ref]

- Waghmare, S. G., Sindhumole, P., Shylaja, M. R., Mathew, D., Francies, R. M., Abida, P. S. and Sajini, S. 2018. Analysis of simple sequence repeat (SSR) polymorphism between N22 and Uma rice varieties for marker assisted selection. *Electronic Journal of Plant Breeding*, **9**(2):511-517. [[Cross Ref](#)]
- Wei, H., Liu, J., Wang, Y., Huang, N., Zhang, X., Wang, L., Zhang, J., Tu, J. and Zhong, X. 2013. A dominant major locus in chromosome 9 of rice (*Oryza sativa* L.) confers tolerance to 48°C high temperature at seedling stage. *Journal of Heredity*, **104**(2):287-294. [[Cross Ref](#)]
- Xiao, Y., Pan, Y., Luo, L., Zhang, G., Deng, H., Dai, L., Liu, X., Tang, W., Chen, L. and Wang, G.L. 2011. Quantitative trait loci associated with seed set under high temperature stress at the flowering stage in rice (*Oryza sativa* L.). *Euphytica*, **178**(3):331-338. [[Cross Ref](#)]
- Xu, Y., Chu, C. and Yao, S. 2021. The impact of high-temperature stress on rice: Challenges and solutions. *The Crop Journal*, **9**(5):963-976. [[Cross Ref](#)]
- Zhang, G., Chen, L., Xiao, G., Xiao, Y., Chen, X. and Zhang, S. 2009. Bulked segregant analysis to detect QTL related to heat tolerance in rice using SSR markers. *Agricultural Sciences in China*, **8**(4):482-487. [[Cross Ref](#)]