



Research Note

Combining ability for yield and yield-associated traits in wheat (*Triticum aestivum* L.)

Afsar Ahmad¹ and Rajesh Kumar Gupta²

Department of Genetics and Plant breeding, Janta Vedic College Baraut Baghpat, Uttar Pradesh – 250611.

E-Mail: afsarahmad06@gmail.com

Abstract

The current breeding effort aims to develop new wheat genotypes with higher genetic potential, thereby contributing to food security. Ten wheat genotypes were crossed in a half-diallel mating design. The F₁ progeny from the 45 crosses, along with their ten parents, were evaluated in a randomized block design to estimate combining ability and the nature of gene action. Analysis of variance for combining ability revealed significant mean squares for both general and specific combining abilities across all studied traits. This indicates the importance of both additive and non-additive gene effects in the inheritance of these traits. The GCA/SCA ratio suggested the predominance of non-additive genetic action for all studied traits. Only five parents (HD3086, UP 2748, WH 1105, UP 2565, and HD 2967) exhibited good general combining ability for yield and associated traits. However, but none of them were found good combiners for all traits. Twenty-one crosses exhibited significant specific combining ability for yield and associated traits. The crosses UP 2748 x HD 3059, UP2565 x HD3086, WH 1105 x UP 2526, WH 1105 x WH 1021, and HD 2967 x WH 1021 were identified as good specific combiners for higher grain yield and associated traits. Utilizing these crosses in breeding programs that exploit heterosis can lead to the creation of new wheat genotypes with a broader genetic base. This approach, which involves crossing high-performing parents for some traits with low-performing parents for others, has the potential to produce offspring with superior characteristics in the next generation due to interactions between dominant genes.

Keyword: General combining ability, Specific combining ability, half diallel, and Wheat.

Wheat (*Triticum aestivum* L.) is a major cereal crop, occupying the largest cultivated area globally. In some Asian countries, wheat is grown for both grain and straw (Beres *et al.*, 2020). It serves as a versatile ingredient in various food products like bread, pasta, and baked goods. Additionally, it has industrial uses in animal feed and biofuels production (Ficco & Borrelli, 2023). The demand for processed wheat products is expected to rise significantly due to population growth, rapid urbanization, and evolving lifestyles (Anonymous, 2015). However, projections indicate a potential decline of 5 to 6 million hectares in wheat cultivation by 2050 due to factors like off-farm diversification and competition from alternative crops (Chaudhary *et al.*, 2023). Combining ability analysis is a widely used technique in wheat breeding. It helps identify promising parental genotypes

and their hybrids by evaluating parental performance in hybrid combinations. This method provides insights into the genetic mechanisms involving numerous major and minor genes that influence wheat yield and related traits (Din *et al.*, 2020).

Some of the major challenges for future wheat production in India include the increasing impact of climate change, particularly rising temperatures and water scarcity, the emergence of new and potentially devastating crop diseases, maintaining sustainable agricultural practices, including reducing dependence on the continuous rice-wheat cropping system, meeting the demands of a growing population with changing urbanization patterns and a preference for higher quality wheat. Despite these challenges, wheat breeders have opportunities to

improve yields by eliminating yield-limiting factors and enhancing the inherent yield potential of wheat varieties (Kumar *et al.*, 2022). Identifying suitable parents for high-yielding breeding programs is crucial. Diallel breeding methods, such as half-diallel crosses, are valuable tools for evaluating general and specific combining abilities for desired traits (Bao *et al.*, 2009). These methods enable breeders to study genetic variation and inheritance patterns in the hybrid offspring. The present study aims to investigate the combining ability of different wheat genotypes and their crosses with respect to yield and yield-related traits using a half-diallel breeding approach.

The current experiment aimed to investigate the genetic basis of yield and associated traits in ten bread wheat genotypes. These genotypes were: HD 2967, WH 1105, UP 2748, UP 2565, UP 2572, UP 2628, UP 2526, HD 3059, HD 3086, and WH 1021. Selection criteria for these genotypes included morphological characteristics, maturity time, and yield-contributing traits. The experiment was conducted at the research farm of Janta Vedic College in Baraut (Baghpat), Uttar Pradesh, India. During the Rabi season of 2018-19, a half-diallel mating design was employed. Ten wheat lines were crossed to generate 45 hybrid combinations. The resulting F_1 progenies, along with their parental lines, were evaluated during Rabi 2019-20 in a randomized block design with three replicates. Each plot consisted of three rows, each one meter long, with a spacing of 22.5 cm between rows and 10 cm between plants. Data on 11 yield and yield-related namely days to heading, days to maturity, flag leaf area (cm^2), plant height at maturity (cm), number of productive tillers per plant, spike length (cm), number of grains per spike, 1000-grain weight, biological yield per plant, grain yield per plant, and harvest index (HI) were collected. For each entry (genotype or cross), five competitive plants were randomly selected, and their average values for all traits (except days to heading and days to maturity) were used for analysis. Data for days to heading and days to maturity were recorded for the entire plot. Combining

ability analysis was performed using Griffing's Method 2, Model I (Fixed) (1956).

This study highlights the importance of understanding both general combining ability (GCA) of parents and specific combining ability (SCA) of progeny for developing high-yielding bread wheat varieties. Analysis of variance between parents, hybrids, and parents versus hybrids revealed the significance of all F_1 traits. As shown in **Table 1**, the mean square for all traits was significantly influenced by both GCA and SCA. This suggests the importance of both additive and non-additive gene action in the inheritance of these traits. The estimated value of SCA variance (σ^2_s) was higher than GCA variance (σ^2_g) for all traits, indicating the predominance of non-additive genes. The ratio of $\sigma^2_{\text{GCA}}/\sigma^2_{\text{SCA}}$ was less than one for all studied traits, further supporting the major role of non-additive gene action. These findings, consistent with previous reports by Singh *et al.* (2012), Nawaz *et al.* (2015), Kumar *et al.* (2017), Gautam *et al.* (2016), Arya *et al.* (2018), Patel (2017), Nassar *et al.* (2020) and Haridy *et al.* (2021) imply that non-additive gene action significantly contributes to the inheritance of all studied traits.

Analysis of GCA for yield and yield-related characteristics (**Table 2**) revealed that only five out of the 10 parents (HD3086, UP 2748, WH 1105, UP 2565, and HD 2967) were good general combiners but none of them were good combiners for all traits. However, some parents showed promise for specific traits: UP2748 and HD 3086 for early flowering and maturity, UP2748 and UP2565 for short stature, WH 1105 and UP2748 for more productive tillers and flag leaf area, UP2748 and HD 3086 for spike length, WH 1105 and HD 3086 for more grains per spike, UP2565 and HD 3086 for 1000-grain weight, HD 2967 and HD 3086 for biomass production, UP2748 and HD 3086 for harvest index and grain yield. These parents (HD3086, UP 2748, WH 1105, UP 2565, and HD 2967) with favorable GCA effects for

Table 1 Analysis of variance (ANOVA) of combining ability analysis for grain yield and yield contributing traits in bread wheat.

Source of variation	d.f.	Mean squares										
		Days to heading	Days to maturity	Plant height (cm)	Productive tillers/plants	Flag leaf Area (cm)	Spike length (cm)	Grains per spike	1000-grain weight	Biological yield/Plant	Harvest Index	Grain yield/Plants
GCA	9	36.68**	11.78**	152.97**	3.97**	70.52**	2.20*	183.37**	18.87**	138.97**	47.31**	67.15**
SCA	45	5.76**	9.06**	31.86**	2.38**	27.61**	2.09**	59.65**	16.83**	164.88**	52.44**	56.25**
Error	108	0.495	0.959	0.236	0.128	0.261	0.038	0.471	0.063	0.351	0.936	0.152
Σg_i^2		1.809	0.541	7.637	0.192	3.513	0.108	9.145	0.940	6.931	2.319	3.350
Σs_{ij}^2		5.262	8.104	31.625	2.255	27.345	2.052	59.183	16.772	164.528	51.500	56.093
$\Sigma g_i^2 / \Sigma s_{ij}^2$		0.344	0.067	0.241	0.085	0.128	0.053	0.155	0.056	0.042	0.045	0.060

*, ** Significant at 0.05 and 0.01 levels of probability, respectively

Table 2. Evaluation of overall General Combining Ability (GCA) effects and performance, including a ranking (of desired traits only) for parental contributions to yield

Parents	Days to heading		Days to maturity		Plant height (cm)		Productive tillers/ plants		Flag leaf Area (cm)		Spike length (cm)	
	Gca	Per se	Gca	Per se	Gca	Per se	gca	Per se	Gca	Per se	gca	Per se
HD- 2967	1.11 **	86.33	0.76 **	125.33	-0.30 **	98.06	0.15 **	7.60	-1.35 **	27.89	-0.20 **	10.89
UP2748	-1.51 **	81.67	-0.52 **	124.33	-3.69 **	84.80	0.36 **	6.60	1.94 **	25.32	0.32 **	11.47
WH 1105	0.02 ns	83.67	0.48 **	124.67	-0.97 **	91.37	0.39 **	7.53	2.28 **	29.82	0.23 **	11.44
UP2565	-0.59 **	82.33	-0.24 ns	123.67	-1.19 **	90.27	0.07 ns	6.40	0.51 **	25.03	-0.03 ns	10.49
UP2572	1.58 **	87.67	0.64 **	125.33	3.55 **	100.23	-0.36 **	7.63	-1.24 **	23.83	-0.40 **	10.87
UP 2628	0.88 **	85.67	-0.36 *	123.33	-0.31 **	98.31	0.09 ns	7.80	-0.92 **	27.67	-0.02 ns	12.77
UP2526	0.19 ns	85.67	-0.08 ns	125.33	0.17 *	90.53	-0.32 **	6.70	0.74 **	28.40	0.06 *	11.80
HD3059	0.08 ns	85.33	0.31 *	125.00	0.67 **	93.74	-0.54 **	5.65	-1.90 **	23.14	-0.30 **	11.30
WH1021	-0.39 **	84.33	0.09 ns	125.00	2.88 **	99.07	-0.19 **	6.50	-0.11 ns	26.24	0.03 ns	12.11
HD3086	-1.37 **	79.67	-1.08 **	119.33	-0.81 **	85.47	0.35 **	8.83	0.04 ns	29.85	0.31 **	12.90
Good general combiner	UP2748, HD 3086, UP 2565, WH 1021		HD 3086, UP2748, UP2628		UP2748,UP2565, WH 1105, HD 3086, UP 2628, HD 2967		WH 1105, UP2748, HD 3086, HD 2967		WH 1105, UP2748, UP 2565		UP2748, HD 3086, WH 1105	
High per se performance	HD3086, UP 2748, UP 2565, WH 1021		HD3086, UP 2748,		UP2748, HD 3086,UP2565, WH 1105, UP 2628, HD 2967		HD 3086, HD 2967, WH 1105, UP2748,		WH 1105, UP2748, UP 2565		HD3086, UP2748, WH 1105	
Common parent	HD3086, WH 1021		HD3086		UP2748, UP 2565, WH 1105, UP 2628, HD 2967		WH 1105, HD 3086, HD 2967, UP 2748		WH 105, UP2748, UP 2565		HD3086, UP2748, WH 1105	

Parents	Grains per spike		1000-grain weight		Biological yield/ Plant		Harvest Index		Grain yield/ Plants	
	Gca	Per se	gca	Per se	Gca	Per se	Gca	Per se	Gca	Per se
HD 2967	0.79 **	53.67	-0.60 **	41.71	3.12 **	64.90	-0.59 **	44.07	0.78 **	28.60
UP2748	-0.21 ns	40.43	0.52 **	38.33	0.79 **	43.26	1.22 **	42.30	1.36 **	18.30
WH 1105	3.74 **	57.73	0.10 *	39.30	1.85 **	56.60	0.28 ns	43.29	1.27 **	24.50
UP2565	0.49 **	44.64	1.35 **	41.40	0.48 **	46.70	0.97 **	41.78	1.05 **	19.51
UP2572	-2.00 **	48.60	-0.23 **	44.23	-1.82 **	54.67	-1.58 **	39.67	-1.64 **	21.68
UP 2628	-1.51 **	45.10	0.34 **	42.30	-0.00 ns	58.32	-1.04 **	47.29	-0.30 **	27.58
UP2526	-0.64 **	48.10	-0.80 **	39.20	-1.31 **	50.00	-0.39 *	41.40	-0.94 **	20.51
HD3059	-2.65 **	42.30	-0.75 **	38.30	-1.88 **	48.23	-1.26 **	39.24	-1.94 **	18.83
WH1021	-1.77 **	39.37	-0.63 **	38.42	-3.00 **	48.01	0.57 **	40.62	-1.26 **	19.50
HD3086	3.75 **	56.20	0.69 **	42.17	1.78 **	65.60	1.82 **	44.46	1.62 **	29.17
Good general combiner	WH 1105, HD 3086, UP 2565		UP2565, HD 3086, UP2748, UP 2628, WH 1105		HD 2967, WH 1105, HD 3086, UP 2748,UP 2565		HD 3086, UP2748, UP 2565, WH 1021		HD 3086, UP2748, WH 1105,UP 2565, HD 2967	
High per se performance	WH 1105, HD3086, HD 2967		UP 2572, UP 2628, HD 2967, UP 2565,		HD 3086, HD 2967, UP 2628, WH 1105, UP 2572		UP 2628, HD 3086, HD 2967, WH 1105		HD 3086, HD 2967, UP 2628, WH 1105, UP 2572	
Common parent based on GCA	WH 1105, HD 3086		UP2565, UP 2628		HD 3086, HD 2967		HD 3086		HD 3086, WH 1105	

*, ** Significant at 0.05 and 0.01 levels of probability, respectively

grain yield and associated traits are considered good general combiners and could be used in future breeding programs to improve grain yield. Based on the overall study, parent HD 3086 exhibited superior performance, making it suitable for recombination programs to produce superior offspring. Similar findings were reported earlier by Singh *et al.* (2013), Singh *et al.* (2014), Kumar *et al.* (2015), Nagar *et al.* (2018), Kumar *et al.* (2017), Hammam *et al.* (2020) and Tariq and Al-Jubouri (2023).

Since SCA effects are often associated with non-additive genes, they are less relevant in self-pollinated crops like wheat because complementary or linkage effects cannot be easily fixed in pure lines. However, SCA can be beneficial for producing transgressive segregants, which are offspring with traits exceeding those of the parents. SCA analysis of 45 crosses revealed that 21 hybrids exhibited significant SCA effects on grain yield and other associated traits (Table 3). The combinations UP 2748 x HD 3059, UP 2565 x HD 3086, WH 1105 x UP 2526, WH 1105 x WH 1021, UP 2748 x UP 2628, UP 2748 x UP 2572, UP 2748 x UP 2565, HD 3059 x WH 1021, UP 2748 x HD 3086, HD 2967 x UP 2572, UP 2565 x WH 1021, UP 2565 x UP 2572, WH 1105 x UP 2628, WH 1105 x HD

3086, UP 2526 x HD 3059, UP 2628 x UP 2526, WH 1105 x UP2565, HD- 2967 x HD3059, UP2748 x UP2526, and UP 2565 x UP 2526 (Table 4) were identified as superior combinations for grain yield and yield associated traits but none of the crosses consistently showed better SCA effect for all traits. Notably, the cross UP 2748 x HD 3086 was the best cross for days to heading, HD 2967 x WH 1105 for days to maturity, HD 2967 x UP 2572 for plant height, HD 2967 x WH 1021 for productive tillers per plant, UP 2565 x UP 2526 for flag leaf area, UP 2748 x UP 2572 for spike length, WH 1105 x UP 2526 for grains per spike, UP 2748 x UP 2572 for thousand grain weight, UP 2748 x HD 3059 for biological yield per plant and UP 2572 x WH 1021 for harvest index. Interestingly, most successful crosses involved parents with contrasting GCA effects (high x low), except for UP 2565 x HD 3086, UP 2748 x UP 2565, and WH 1105 x HD 3086 (high x high). This suggested that high specific combining ability was not always linked to high parental GCA effects, potentially due to the importance of intrallelic interaction or additive x additive gene action in high x high combinations.

The current study demonstrates that high x high GCA parents may not always produce the best recombinants.

Table 3 Best cross combinations with significant and desirable SCA effects for yield and yield-associated traits

S. No.	Crosses	SCA effects	GCA effects		GCA effects of parent	Nature of gene of action	Yield contributing traits
			P ₁	P ₂			
1	UP 2748 x HD 3059	7.86	1.36	-1.94	H x L	Non-additive	DH, PT, FLA, SL, GS, TGW, BY
2	UP 2565 x HD 3086	7.63	1.05	1.62	H x H	Additive	DH, DM, PT, FLA, SL, GS, TGW, BY, HI
3	WH 1105 x UP 2526	6.99	1.27	-0.94	H x L	Non-additive	PH, PT, FLA, SL, GS, TGW, BY, HI,
4	WH 1105 x WH 1021	6.87	1.27	-1.26	H x L	Non-additive	PH, PT, FLA, SL, GS, TGW, BY, HI
5	HD 2967 x WH 1021	6.75	0.78	-1.26	H x L	Non-additive	DH, PT, FLA, SL GS, BY, HI
6	UP 2748 x UP 2628	4.91	1.36	-.30	H x L	Non-additive	DH, DM, PH, FLA, PT, TGW, SL, GS, BY, HI,
7	UP 2748 x UP 2572	4.11	1.36	-1.64	H x L	Non-additive	PH, FLA, PT, TGW, SL, GS, BY, HI,
8	UP 2748 x UP 2565	4.16	1.36	1.05	H x H	Additive	PH, FLA, SL, GS, BY, HI
9	HD 3059 x WH 1021	3.42	-1.94	-1.26	L x L	Non-additive	PH, PT, SL, GS, TGW, BY,
10	UP 2748 x HD 3086	3.26	1.36	1.62	H x L	Non-additive	DH, PT, FLA, SL, GS, TGW, BY
11	HD 2967 x UP 2572	3.02	0.78	-1.64	H x L	Non-additive	DM, PH, PT, FLA, SL, TGW, HI,
12	UP 2565 x WH 1021	2.70	1.05	-1.26	L x L	Non-additive	DM, PH, PT, GS, TGW, BY, HI
13	UP 2565 x UP 2572	2.13	1.05	-1.64	H x L	Non-additive	PH, PT, FLA, SL, TGW, BY, HI
14	WH 1105 x UP 2628	2.13	1.27	-.30	H x L	Non-additive	PH, PT, FLA, SL
15	WH 1105 x HD 3086	1.48	1.27	1.62	H x H	Additive	PT, SL, GS, TGW, HI
16	UP 2526 x HD 3059	1.15	-0.94	-1.94	L x L	Non-additive	DH, DM, PT, GS, TGW, BY,
17	UP 2628 x UP 2526	0.63	-0.30	-0.94	L x L	Non-additive	DH, PH, PT, FLA, SL
18	WH 1105 x UP2565	0.44	1.27	1.05	H x H	Additive	DH, SL, GS, TGW, HI
19	HD- 2967 X HD3059	0.41	0.78	-1.94	H x L	Non-additive	DH, FLA, SL, GS, HI,
20	UP2748 X UP2526	0.21	1.36	-0.94	H x L	Non-additive	DM, PT, FLA, TGW,
21	UP 2565 x UP 2526	0.22	1.05	-0.94	H x L	Non-additive	PH, FLA, BY,

DH= Days to heading, DM = Day to maturity, PH = Plant height, FLA = Flag leaf area, PT = Productive tillers per plant, SL = Spike length, GS= Grains per spike, TGW = Thousand-grain weight, BY = Biological yield per plant, HI = harvest index.

Table 4. Elite cross combinations for each trait identified based on SCA effects

Character	Cross combination	SCA effects	GCA effects		GCA effects on parents	Nature of gene action
			P ₁	P ₂		
Days to heading	UP 2748 x HD 3086	-1.74	-1.51	-1.37	H x H	Additive
Days to maturity	HD 2967 x WH 1105	-3.92	0.76	0.48	L x H	Non-additive
Plant height (cm)	HD 2967 x UP 2572	-8.55	-0.30	3.55	H x L	Non-additive
Productive tillers per plant	HD 2967 x WH 1021	2.15	0.15	-0.19	H x L	Non-additive
Flag leaf area (cm)	UP 2565 x UP 2526	7.63	0.51	0.74	H x H	Additive
Spike length (cm)	UP 2748 x UP 2572	1.38	0.32	-0.40	H x L	Non-additive
Grains per spike	WH 1105 x UP 2526	7.80	3.74	-0.64	H x L	Non-additive
1000-grain weight	UP 2748 x UP 2572	3.87	0.52	-0.23	H x L	Non-additive
Biological yield per plant	UP 2748 x HD 3059	18.56	0.79	-1.88	H x L	Non-additive
Harvest index	UP 2572 x WH 1021	7.73	-1.58	0.57	L x L	Non-additive
Grain yield per plant	UP 2748 x HD 3059	7.86	1.36	-1.94	H x L	Non-additive

Conversely, crosses involving parents with contrasting GCA effects (high x low) can generate transgressive segregants due to dominance gene interactions. These findings align with previous research by Singh *et al.* (2007), Singh *et al.* (2012), Singh *et al.* (2013), Rahul *et al.* (2017), Kumar *et al.* (2017), Rania A. R. El-Said (2018), Khokhar *et al.* (2019), Nagar *et al.* (2020), Joshi *et al.* (2020) and Bilgin *et al.*, (2022).

The study identified five parents-HD3086, UP2748, WH1105, UP2565, and HD2967 – as promising due to their good general combining ability for yield and related traits. These parents can be directly used in breeding programs to enhance grain yield in wheat. Furthermore, specific crosses, such as UP2748 x HD3059 and UP 2565 x HD 3086, showed significant potential for improving grain yield and associated characteristics in future bread wheat breeding efforts. These promising crosses likely benefit from both additive and non-additive gene effects. Breeding approaches like bi-parental mating and diallel selective mating could be particularly effective in improving bread wheat characteristics. These methods allow for crosses between selected parents in multiple cycles, maximizing the utilization of both additive and non-additive genetic effects.

REFERENCES

- Anonymous. 2015. Progress Report vision 2050 of All India Coordinated Wheat and Barley Improvement Project 2015, Project Director's Report. Ed: Indu Sharma. Directorate of Wheat Research, Karnal, India. pp. 6
- Arya, V.K., Kumar, P., Singh, J., Kumar, L. and Sharma, A.K. 2018. Genetic analysis of some yield and quality traits in bread wheat (*Triticum aestivum* L.). *Wheat and Barley Research*, **10**(1): 25-32. [Cross Ref]
- Bao, Y., Wang, S., Wang, X., 2009. Heterosis and combining ability for major yield traits of a new wheat germplasm Shannong 0095 derived from *Thinopyrum intermedium*. *Agricultural Sciences in China*, **8**(6):753–760. [Cross Ref]
- Beres, B.L., Rahmani, E., Clarke, J.M., Grassini, P., Pozniak, C.J., Geddes, C.M., Porker, K.D., May, W.E. and Ransom, J.K. 2020 A systematic review of durum wheat: Enhancing production systems by exploring genotype, environment, and management (G × E × M) synergies. *Front. Plant Sci.* **11**:568657. [Cross Ref]
- Bilgin, O., Yazici, E., Balkan, A. and Başer, I. 2022. Selection for high yield and quality in half-diallel bread wheat F₂ populations (*Triticum aestivum* L.) through heterosis and combining ability analysis. *International Journal of Agriculture Environment and Food Sciences*, **6**(2): 285-293. [Cross Ref]
- Chaudhary Divya, Swati, Kumar Anil, Reddy Basu Sudhakar and Rohit. 2023. Genetic variability and character association studies in bread wheat *Triticum aestivum* L. under two different water regimes. *Electronic Journal of Plant Breeding*, **14**(4): 1330-1336. [Cross Ref]
- Din, K., Khan, N. U., Gul, S., Khan, S. U., Khalil, I. H., Khan, S. A. and Khalil, I. A. 2021. Line by tester combining ability analysis for earliness and yield traits in bread wheat (*Triticum aestivum* L.). *Japs: Journal of animal & plant sciences*, **31**(2). [Cross Ref]
- Ficco, D. B. M. and Borrelli, G. M. 2023. Nutritional Components of Wheat Based Food: Composition, Properties, and Uses. *Foods*, **12**(21):4010. [Cross Ref]
- Gautam, A., Sai Prasad, S.V., Ambati, D. and Jajoo, A. 2016. Genetic analysis of grain yield and its contributing components in diallel crosses of durum wheat

- (*Triticum durum* L.) under terminal heat stress conditions. *Journal of Wheat Research*, **8**(2): 12-18.
- Hammam, O. Kh., abd-El Zaher, I.N., Haridy, M.H. and Al-Aref Kh. A.I., 2020. Estimate of combining ability and correlation for yield and its components in bread wheat (*Triticum aestivum* L.). *Archives of Agriculture science journal*, **3**(3): 1-12.
- Haridy, M. H., El-Zaher, A. and Mahdy, A. Y. 2021. Estimate of combining ability and correlation for some bread wheat genotypes. *Assiut Journal of Agricultural Sciences*, **52**(4): 1-11. [Cross Ref]
- Joshi, A., Kumar, A. and Kashyap, S. 2020. Genetic analysis of yield and yield contributing traits in bread wheat. *IJAEB*, **13**(2): 119-128. [Cross Ref]
- Khokhar, A.A., Nizamani, F.G., Rind, R.A., Nizamani, M.M., Khokhar, M.U., Shah, A., Nizamani, A.L. and Rind, M.R. 2019. Combining ability estimates in 6 x 6 half diallel crosses of bread wheat (*Triticum aestivum* L.). *Pure Appl. Biol.*, **8**(3): 1980-1990. [Cross Ref]
- Kumar, B., Bharadwaj, D.N., Singh, L. and Gupta, S.K. 2017. Estimation of combining ability in F_1 and F_2 generations of diallel crosses in winter wheat (*Triticum aestivum* L. Em. Thell). *Int. J. Curr. Microbiology App. Sci*, **6**(12): 1432-1444. [Cross Ref]
- Kumar, J., Singh, S.K., Singh, L., Kumar, M., Srivastava, M., Singh, J. and Kumar, A. 2017. Combining ability analysis for yield and its components in bread wheat (*Triticum aestivum* L.) under abiotic stress. *Int. J. Curr. Microbiol. App. Sci.*, **6**(3): 24-39. [Cross Ref]
- Kumar, L., Chhogyel, N., Gopalakrishnan, T., Hasan, M. K., Jayasinghe, S. L., Kariyawasam, C. S., ... and Ratnayake, S. 2022. Climate change and future of agri-food production. In *Future foods* (pp. 49-79). Academic Press. [Cross Ref]
- Kumar, P., Singh, G., Singh, Y. P., Abhishek, D. and Nagar, S.S. 2015. Study of combining ability analysis in half-diallel crosses of spring wheat (*Triticum aestivum* L.). *International Journal of Advanced Research*, **3**(9): 1363-1370.
- Nagar, S. S., Kumar, P., Vishwakarma, S.R. and Gupta, V. 2018. Diallel analysis of some grain yield traits in wheat. *Wheat and Barley Research*, **10**(1): 45-51. [Cross Ref]
- Nagar, S.S., Kumar, P., Singh, G., Gupta, V., Singh, C. and Tyagi, B.S. 2020. Assessing gene action utilizing Hayman's graphical approach in bread wheat (*Triticum aestivum* L.). *Journal of Crop and Weed*, **16**(1): 29-37. [Cross Ref]
- Nassar, M. A. A., Elbanna, M. N., Elsorady, G. A.I. and Nofal, M. E. A. 2020. Yield and its components of some half diallel bread wheat crosses. *Egypt. Acad. J. Biolog. Sci.*, **11**(2): 21-33. [Cross Ref]
- Nawaz, M.S., Ansari, B.A., Kumbhar, M.B. and Keerio, M.I. 2015. Combining ability estimates of parents and F_1 hybrids for economically useful traits of hexaploid wheat (*Triticum*
- Patel, H.N. 2017. Combining ability analysis for yield and its components in bread wheat. *Electronic Journal of Plant Breeding*, **8**(2): 404-408. [Cross Ref]
- Rahul Bhardwaj, R.S. Sain, Rajdeep Mundiyyara, and Sharma Ankita. 2017. Combining ability analysis of F_1 and F_2 generation of ten parent half diallel in bread wheat (*Triticum Aestivum* L. Em. Thell). *International Journal of Agriculture Sciences*, **9**(22): 4238-4241
- Rania A. R. El-Said. 2018. Assessment of genetical parameters of yield and its attributes in bread wheat (*Triticum aestivum*, L.). *J. Agric. Chem. and Biotechn. Mansoura Univ.*, **9** (10): 243-251. [Cross Ref]
- Singh Rajesh, Pandey H.N., Mishra M. M., Sai Prasad S. V. and Parikh Mangla. 2007. Combining ability and heterosis for yield and grain quality in durum wheat (*Triticum turgidum* var. durum) *Madras Agriculture Journals*, **94**: (1-6)
- Singh, K., Singh, U.B and Sharma, S.N. 2013. Combining ability analysis for yield and its components in bread wheat (*Triticum aestivum* L. em. Thell.). *J. Wheat Res.* **5** (1), 63-67.
- Singh, V., Krishna, R., Singh, S. and Vikram, P. 2012. Combining ability and heterosis analysis for yield traits in bread wheat (*Triticum aestivum*). *Indian Journal of Agricultural Sciences*, **82**(11): 916-21. [Cross Ref]
- Tariq, R. and Al-Jubouri, J. M. (2023, December). Combining Ability and Gene Action of Half Diallel Crosses in Bread Wheat (*Triticum aestivum* L.). In *IOP Conference Series: Earth and Environmental Science* (Vol. 1262, No. 5, p. 052027). IOP Publishing. [Cross Ref]