


# Electronic Journal of Plant Breeding



## Research Note

### Inter and intra cross variability analysis in F<sub>3</sub> generation of mung bean [*Vigna radiata* (L.) Wilczek]

P. K. Nayak, C.L. Khatik\*, M. Khan, Kailash Chandra, J. Lal, D. Tripathi, K.C. Verma and M.A. Khan

Department of GPB, SKNCOA, Jobner and Agricultural Research Station, Fatehpur-Shekhawati, Sikar (SKNAU, Jobner) Rajasthan-India

\*E-Mail: champalalkhatik@gmail.com

#### Abstract

The present investigation was conducted to evaluate 12 crosses of mungbean genotypes for 11 characters to access the extent of genetic variability and to comprehend the heritable aspect of variation for grain yield and its related characters. All the 12 crosses were raised in a compact family block design with three replications in two environments during the *kharif* season of 2022. The analysis of variance indicated significant variations among the families for grain yield and all other characters in both environments. Conversely, differences among the progenies within a family varied depending on specific trait and environment (E1) to environment (E2). Inter cross variability revealed high GCV and PCV for primary branches per plant in both environments. High estimates of PCV were recorded for pods per plant and grain yield per plant in both environments and for clusters per plant in E2. High estimates of heritability were exhibited by primary branches per plant in both the environments and plant height in E1. High estimates of genetic advance as per cent of mean were observed for primary branches per plant and plant height in both environments. Inter cross variability revealed that the highest magnitude of heritability was recorded by family C1 and C7 for grain yield per plant in E1 and E2 respectively. The highest magnitude of genetic advance as per cent of mean was exhibited by family C3 for primary branches per plant in E1 and by family C12 for clusters per plant in E2.

**Keywords:** Genetic variability, Inter cross variability, Heritability, Mungbean

Mungbean is a traditional and well-known tropical leguminous crop of Asia because of its nutritional value and cropping system adaptability. Mungbean, also known as moong, is the third most significant legume, next to chickpea and pigeon pea. According to Karpechenko (1925), it bears the diploid chromosomal number  $2n = 2x = 22$ . According to Vavilov (1926) mungbean is a native of India and Central Asia. Since prehistoric times, it has been grown in these regions. The diploid mungbean ( $2n = 22$ ) is a member of the Leguminaceae family, Papilionaceae subfamily, genus *Vigna* and species *radiata*. About 18.0 to 32.0 per cent protein and 1 to 5 per cent fat are both found in pulses. Chemical components include proteins, fatty acids, carbohydrates, vitamin B1, vitamin B2, beta-carotene, folic acid, calcium,

phosphorus, and iron. India accounts for around 25 per cent of global production from 35 per cent of global area under pulses. In India mungbean occupied roughly 31.15 lakh hectares during 2019–20 with a production of 23.40 lakh tonnes and productivity of 798 kg/ha. States that produce a significant amount of mungbean in India include Rajasthan, Maharashtra, Karnataka, Madhya Pradesh, Odisha and Telangana (Anonymous, 2019-20). In Rajasthan, the total area planted with mungbean during 2019–2020 was 23.27 lakh hectares, with a total production of 13.04 lakh tonnes and a productivity of 561 kg/ha and it is the leading mung bean producing state in the country (Anonymous, 2019-20). It is primarily grown in the semi-arid and dry regions of Rajasthan, particularly in the districts of Nagaur, Ajmer, Jaipur, Jodhpur, Jalore,

Table 1. Details of mung bean research materials

crosses	F <sub>2</sub> plants selected	Codes in F <sub>3</sub> 's	
		Family	Progeny
HUM 1 × IPM-205-07 (A <sub>1</sub> ) (A <sub>2</sub> )	A <sub>1</sub> A <sub>2</sub> -1 A <sub>1</sub> A <sub>2</sub> -2 A <sub>1</sub> A <sub>2</sub> -3 A <sub>1</sub> A <sub>2</sub> -4 A <sub>1</sub> A <sub>2</sub> -5	C <sub>1</sub>	C1P1 C1P2 C1P3 C1P4 C1P5
HUM 1 × MH 421 (A <sub>1</sub> ) (A <sub>3</sub> )	A <sub>1</sub> A <sub>3</sub> -1 A <sub>1</sub> A <sub>3</sub> -2 A <sub>1</sub> A <sub>3</sub> -3 A <sub>1</sub> A <sub>3</sub> -4 A <sub>1</sub> A <sub>3</sub> -5	C <sub>2</sub>	C2P1 C2P2 C2P3 C2P4 C2P5
HUM 1 × LGG 460 (A <sub>1</sub> ) (A <sub>4</sub> )	A <sub>1</sub> A <sub>4</sub> -1 A <sub>1</sub> A <sub>4</sub> -2 A <sub>1</sub> A <sub>4</sub> -3 A <sub>1</sub> A <sub>4</sub> -4 A <sub>1</sub> A <sub>4</sub> -5	C <sub>3</sub>	C3P1 C3P2 C3P3 C3P4 C3P5
HUM 1 × MH-2-15 (A <sub>1</sub> ) (A <sub>5</sub> )	A <sub>1</sub> A <sub>5</sub> -1 A <sub>1</sub> A <sub>5</sub> -2 A <sub>1</sub> A <sub>5</sub> -3 A <sub>1</sub> A <sub>5</sub> -4 A <sub>1</sub> A <sub>5</sub> -5	C <sub>4</sub>	C4P1 C4P2 C4P3 C4P4 C4P5
HUM 1 × RMG-975 (A <sub>1</sub> ) (A <sub>6</sub> )	A <sub>1</sub> A <sub>6</sub> -1 A <sub>1</sub> A <sub>6</sub> -2 A <sub>1</sub> A <sub>6</sub> -3 A <sub>1</sub> A <sub>6</sub> -4 A <sub>1</sub> A <sub>6</sub> -5	C <sub>5</sub>	C5P1 C5P2 C5P3 C5P4 C5P5
IPM-205-07 × MH 421 (A <sub>2</sub> ) (A <sub>3</sub> )	A <sub>2</sub> A <sub>3</sub> -1 A <sub>2</sub> A <sub>3</sub> -2 A <sub>2</sub> A <sub>3</sub> -3 A <sub>2</sub> A <sub>3</sub> -4 A <sub>2</sub> A <sub>3</sub> -5	C <sub>6</sub>	C6P1 C6P2 C6P3 C6P4 C6P5
IPM-205-07 × LGG 460 (A <sub>2</sub> ) (A <sub>4</sub> )	A <sub>2</sub> A <sub>4</sub> -1 A <sub>2</sub> A <sub>4</sub> -2 A <sub>2</sub> A <sub>4</sub> -3 A <sub>2</sub> A <sub>4</sub> -4 A <sub>2</sub> A <sub>4</sub> -5	C <sub>7</sub>	C7P1 C7P2 C7P3 C7P4 C7P5
IPM-205-07 × ML-818 (A <sub>2</sub> ) (A <sub>7</sub> )	A <sub>2</sub> A <sub>7</sub> -1 A <sub>2</sub> A <sub>7</sub> -2 A <sub>2</sub> A <sub>7</sub> -3 A <sub>2</sub> A <sub>7</sub> -4 A <sub>2</sub> A <sub>7</sub> -5	C <sub>8</sub>	C <sub>8</sub> P1 C <sub>8</sub> P2 C <sub>8</sub> P3 C <sub>8</sub> P4 C <sub>8</sub> P5
IPM-205-07 × MH-2-15 (A <sub>2</sub> ) (A <sub>5</sub> )	A <sub>2</sub> A <sub>5</sub> -1 A <sub>2</sub> A <sub>5</sub> -2 A <sub>2</sub> A <sub>5</sub> -3 A <sub>2</sub> A <sub>5</sub> -4 A <sub>2</sub> A <sub>5</sub> -5	C <sub>9</sub>	C <sub>9</sub> P1 C <sub>9</sub> P2 C <sub>9</sub> P3 C <sub>9</sub> P4 C <sub>9</sub> P5
IPM-205-07 × RMG-975 (A <sub>2</sub> ) (A <sub>6</sub> )	A <sub>2</sub> A <sub>6</sub> -1 A <sub>2</sub> A <sub>6</sub> -2 A <sub>2</sub> A <sub>6</sub> -3 A <sub>2</sub> A <sub>6</sub> -4 A <sub>2</sub> A <sub>6</sub> -5	C <sub>10</sub>	C10P1 C10P2 C10P3 C10P4 C10P5
ML-818 × RMG-975 (A <sub>7</sub> ) (A <sub>6</sub> )	A <sub>7</sub> A <sub>6</sub> -1 A <sub>7</sub> A <sub>6</sub> -2 A <sub>7</sub> A <sub>6</sub> -3 A <sub>7</sub> A <sub>6</sub> -4 A <sub>7</sub> A <sub>6</sub> -5	C <sub>11</sub>	C11P1 C11P2 C11P3 C11P4 C11P5
MH-2-15 × RMG-975 (A <sub>5</sub> ) (A <sub>6</sub> )	A <sub>5</sub> A <sub>6</sub> -1 A <sub>5</sub> A <sub>6</sub> -2 A <sub>5</sub> A <sub>6</sub> -3 A <sub>5</sub> A <sub>6</sub> -4 A <sub>5</sub> A <sub>6</sub> -5	C <sub>12</sub>	C12P1 C12P2 C12P3 C12P4 C12P5

Pali, Sikar and Jhunjhunu. Inter cross variability refers to the variation observed among different populations, which are derived from different parental crosses. Intra cross variability, on the other hand, refers to the variation observed within a single population, which is derived from a single parental cross. To genetically modify quantitative traits in any crop, a full understanding of the type and degree of variability present in the crop, as well as the heritability of the desirable attributes inherent in the breeding material, is required. For the establishment of an effective breeding plan, a study of genetic variability using several metrics such as heritability, genetic progress and phenotypic and genotypic coefficient of variation is essential. Understanding distinct genetic parameter estimations is critical for improving crop yield by better using the type and degree of genetic diversity available in the base material.

Plant material: Five single plants with high yield in  $F_2$  generation in each of the 12 crosses were selected and advanced to  $F_3$  generation. Each cross was considered a family, and the selected individual plants within a cross were referred to as progenies. All the twelve crosses were grown in compact family block design with three replications in conducted in two environments namely Experimental Farm of Department of GPB, SKNAU,

Jobner and the ARS, Fatehpur, Shekhawati, Sikar during *Kharif*, 2022. In both the locations, the experimental plot consisted of 3 m long single rows spaced 30 cm apart, with an intra row spacing of 10 cm. Data were recorded on five randomly selected plants in each row for the traits *viz.* plant height, primary branches /plant, clusters /plant, pods/ cluster, pods/ plant, pod length, seeds /pod, 100-seed weight, grain yield /plant. Days to 50 % flowering and days to maturity were recorded on the basis of individual progeny rows.

Statistical analysis: The analysis of variance for between families and between progenies within a family was worked out using the method proposed by Panse and Sukhatme (1978). Genotypic and phenotypic coefficient of variation was estimated as per Burton (1952). Heritability in broad sense was estimated according to Burton and Devane (1953). Genetic advance and genetic advance as per cent of mean were determined as per the formula suggested by Robinson *et al.* (1949) and Johnson *et al.* (1955).

The estimates of genetic variability parameters between the families and within the families were computed for 11 traits in 12 crosses across under two environments are presented in **table 2 and table 3**.

**Table 2. Estimation of variability parameters between families for yield related characters in two environments**

S. No.	Characters	Env	Mean	EV	GV	PV	GCV	PCV	H(bs)	GA	GAM
1	Days to 50 per cent flowering	E1	37.72	1.06	0.37	1.43	1.61	3.17	25.70	0.63	1.68
		E2	37.94	1.09	0.33	1.43	1.52	3.15	23.41	0.58	1.52
2	Days to maturity	E1	69.28	1.95	1.90	3.86	1.99	2.84	49.35	2.00	2.88
		E2	69.63	2.10	0.86	2.96	1.33	2.47	29.07	1.03	1.48
3	Plant height	E1	69.06	38.61	71.58	110.19	12.25	15.20	64.96	14.05	20.34
		E2	45.85	39.52	42.53	82.05	14.22	19.75	51.83	9.67	21.09
4	Primary branches per plant	E1	0.89	0.03	0.14	0.17	41.41	46.29	80.03	0.68	76.31
		E2	1.96	0.05	0.17	0.21	20.92	23.63	78.40	0.75	38.16
5	Clusters per plant	E1	5.25	0.82	0.08	0.90	5.39	18.06	8.90	0.17	3.31
		E2	4.15	0.80	0.62	1.42	18.97	28.74	43.57	1.07	25.79
6	Pods per cluster	E1	3.54	0.37	0.06	0.43	7.05	18.65	14.31	0.19	5.50
		E2	4.38	0.74	0.20	0.94	10.23	22.18	21.27	0.43	9.72
7	Pods per plant	E1	17.45	9.65	4.59	14.24	12.28	21.62	32.25	2.51	14.36
		E2	16.98	7.76	4.33	12.09	12.26	20.48	35.80	2.56	15.10
8	Pod length	E1	7.52	0.57	0.17	0.74	5.42	11.43	22.49	0.40	5.30
		E2	7.46	0.62	0.09	0.71	3.96	11.26	12.36	0.21	2.87
9	Seeds per pod	E1	11.17	1.07	0.10	1.17	2.88	9.68	8.83	0.20	1.76
		E2	11.25	1.03	0.11	1.14	2.92	9.49	9.49	0.21	1.86
10	100-seed weight	E1	3.43	0.11	0.06	0.17	7.06	11.86	35.43	0.30	8.66
		E2	3.37	0.10	0.06	0.16	7.07	11.83	35.65	0.29	8.69
11	Grain yield per plant	E1	4.16	0.67	0.21	0.88	10.96	22.53	23.66	0.46	10.98
		E2	4.45	0.77	0.13	0.90	8.03	21.34	14.17	0.28	6.23

**Table 3. Estimation of variability parameters in intra family progenies having significant difference in E1 and E2 environments for yield**

S. No.	Characters	Env	Families	Mean	EV	GV	PV	GCV	PCV	H(bs)	GA	GAM		
1	Plant height	E1	C5	71.95	23.11	25.94	49.06	7.08	9.73	52.88	7.63	10.60		
			C7	60.65	18.65	26.20	44.86	8.44	11.04	58.42	8.06	13.29		
			C8	60.65	55.23	55.13	110.41	12.24	17.33	49.93	10.80	17.82		
		E2	C1	52.55	37.67	40.07	77.75	12.05	16.78	51.54	9.36	17.81		
			C12	50.30	31.57	50.41	81.92	14.11	17.99	61.53	11.47	22.80		
			C1	1.24	0.07	0.15	0.22	30.93	37.79	66.98	0.65	52.15		
2	Primary branches per plant	E1	C2	0.83	0.04	0.80	0.12	34.21	41.54	67.82	0.048	58.04		
			C3	0.75	0.04	0.08	0.12	38.06	46.21	67.86	0.49	64.59		
			C9	0.91	0.05	0.10	0.16	36.36	43.75	69.06	0.56	62.25		
		E2	C10	1.65	0.15	0.15	0.31	23.64	33.53	49.70	0.57	34.33		
			C1	2.24	0.07	0.15	0.22	17.13	20.93	67.00	0.65	28.89		
			C3	1.75	0.06	0.04	0.10	8.98	14.89	36.39	0.23	11.16		
		3	Clusters per plant	E1	C4	1.81	0.04	0.09	0.12	16.11	19.19	70.44	0.50	27.85
					C7	1.91	0.04	0.109	0.15	17.27	20.77	69.10	0.56	29.57
					C8	2.65	0.15	0.15	0.31	14.72	20.88	49.71	0.57	21.38
				E2	C11	1.91	0.05	0.11	0.16	17.27	20.78	69.10	0.56	29.58
					C12	2.63	0.15	0.15	0.31	14.72	20.88	49.71	0.57	21.38
					C2	5.56	1.18	1.15	2.33	18.96	26.95	49.53	1.56	27.50
4	Pods per cluster	E1	C8	5.29	0.60	0.69	1.30	15.71	21.51	53.33	1.25	23.63		
			C11	4.55	1.09	1.94	3.03	30.62	38.28	64.01	2.29	50.47		
			C4	4.73	0.79	1.19	1.98	23.08	29.78	60.09	1.74	36.86		
		E2	C8	4.52	1.76	1.69	3.45	28.78	41.13	48.95	1.87	41.48		
			C12	3.70	2.30	2.32	4.62	41.22	58.13	50.28	2.23	60.21		
			C5	3.80	0.42	0.71	1.13	22.12	27.91	62.84	1.37	36.13		
5	Grain yield per plant	E1	C6	3.79	0.34	0.37	0.72	16.09	22.29	52.12	0.91	23.93		
			C8	3.62	0.39	0.84	1.23	25.36	30.60	68.67	1.57	43.29		
			C9	3.61	0.51	0.80	1.31	24.76	31.69	61.07	1.44	39.86		
		E2	C10	3.59	0.49	0.47	0.96	19.05	27.27	48.83	0.98	27.42		
			C12	3.17	0.53	0.25	0.78	15.86	27.92	32.27	0.59	18.56		
			C1	4.80	0.21	0.33	0.53	11.88	15.21	61.04	0.92	19.13		
6	Grain yield per plant	E2	C2	4.85	0.36	0.43	0.79	13.48	18.33	54.06	0.99	20.42		
			C4	4.31	0.19	0.38	0.57	14.36	17.59	66.65	1.04	24.15		
			C7	4.30	0.21	0.36	0.58	13.98	17.62	62.96	0.98	22.85		

Genotypic and phenotypic coefficient of variation: Estimates of coefficient of variation indicated that the magnitude for PCV was higher than that of GCV but the difference between phenotypic and genotypic coefficient variation suggests little influence of environment on the expression of characters. High GCV and PCV was observed for primary branches per plant in both the environments. Such results were also reported by Gadakh *et al.* (2013). Moderate GCV and high PCV was recorded for pods per plant in both the environments, clusters per plant in E2, pods per cluster in E2, grain yield per plant in E1. This agrees with the results reported by Narasimhulu *et al.* (2013). Low GCV & moderate PCV were observed

for pod length, 100-seed weight in both the environments, clusters per plant and pods per cluster in E1. This was also reported by Lakshminarayanareddy *et al.* (2003). Days to 50 % flowering, days to maturity, seeds per pod in both the environments exhibited low GCV & PCV, which were in accordance with the findings of Makeen *et al.* (2007), Tabasum *et al.* (2010), Nand and Anuradha (2013), Jeberson *et al.* (2015), Yadav *et al.* (2017), Mehandi *et al.* (2018), Sirohi *et al.* (2018), Susmitha and Jayamani (2018), Anuradha *et al.* (2019), Pavan *et al.* (2019). Low GCV & high PCV was observed only for grain yield per plant in E2. Moderate GCV and PCV were observed for plant height in both the environments. Among the above

discussed superior families in E1, C3, C1, C7 recorded high GCV and PCV, C4 recorded high PCV and moderate GCV while C2 and C11 recorded moderate GCV and PCV for grain yield per plant. However, in E2 C1 and C10 recorded high GCV and PCV, C2, C11 and C12 recorded high PCV and moderate GCV while C7 recorded moderate GCV and PCV. Similar trends were reported by Pawar *et al.* (1988), Kumar *et al.* (1996) and Pawar *et al.* (2001).

Heritability and Genetic advance: High heritability was observed for primary branches per plant in both the environments and plant height in E1. Similar finding was also reported by Gadakh *et al.* (2013). Moderate heritability and high genetic advance were observed for plant height and clusters per plant in E2. Moderate heritability coupled with moderate genetic advance was observed for pods per plant in both the environments. Moderate heritability coupled with low genetic advance was observed for 100-seed weight in both the environments and days to maturity in E1. Similar findings were reported by Kumar *et al.* (2010), Ramachandra and Lavanya (2017), Pavan *et al.* (2019), Yoseph *et al.* (2022) and Prajapati *et al.* (2023). Low heritability coupled with low genetic advance were observed for days to 50 per cent flowering, seeds per pod, pods per cluster and pod length in both the environments, days to maturity and grain yield per plant in E2, clusters per plant in E1. Low heritability coupled with moderate genetic advance were observed only for grain yield per plant in E1.

Among the above discussed superior families in E1 C1 and C2 recorded high heritability while C3, C4, C7 and C11 recorded moderate heritability for grain yield per plant. High genetic advance as per cent of mean was observed for C1, C3, C4 and C7 while C2 and C11 recorded moderate genetic advance as per cent of mean for grain yield per plant. High heritability coupled with high genetic advance was observed for C1. This shows the preponderance of additive gene action and heritability is due to additive gene effect and selection in this family for yield and component traits will be effective. Moderate heritability coupled with high genetic advance was observed for C3, C4 and C7. This indicates the presence of additive gene effect and heritability is due to high environmental influence. Selection within these families may be effective. High heritability coupled with moderate genetic advance was observed for C2. This indicates the presence of non-additive gene effect and heritability is due to unfavourable environmental effect. Simple selection within these families may not be effective. Moderate heritability coupled with moderate genetic advance was observed for C11. This indicates that the character is highly influenced by environmental effects. Selection within this family may not be rewarding.

On the other hand, in E2 C1, C2 and C7 recorded high heritability while C10, C11 and C12 recorded moderate heritability for grain yield per plant. High genetic advance as per cent of mean was observed for C1, C2, C7 and C10

while C11 and C12 recorded moderate genetic advance as per cent of mean for grain yield per plant. High heritability coupled with high genetic advance was observed for C1, C2 and C7. This shows the preponderance of additive gene action and heritability is due to additive gene effect and selection in these families for yield and component traits will be effective. Moderate heritability coupled with high genetic advance was observed for C10. This indicates the presence of additive gene effect and heritability is due to high environmental influence. Selection within these families may be effective. Moderate heritability coupled with moderate genetic advance was observed for C11 and C12. Selection within this family may not be rewarding. These are in accordance with the findings of Idress *et al.* (2006), Narasimhulu *et al.* (2013), Tiwari *et al.* (2014) and Sofia *et al.* (2017). This indicates that the character is highly influenced by environmental effects.

Thus selection in the families C1 and C2 which were found to possess high heritability and genetic advance for grain yield per plant in both the environments could yield desirable results.

## REFERENCES

- Anonymous, 2019-20. Agriculture Statistics. Department of Agriculture. Government of Rajasthan.
- Anuradha, N., Patro, T.S.S.K. and Kumar, R.S. 2019. Study on genetic variability, heritability and correlation analysis for grain yield and yield attributing traits in green gram [*Vigna radiata* (L.) Wilczek]. *IJCS*, **7**(3): 2050-2052.
- Burton, G.W. 1952. Quantitative inheritance in grasses. *Proceedings of the Sixth International Grassland Congress*, **1**: 277-282.
- Burton, G.W. and DeVane, E.W. 1953. Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. *Agro. J.*, **4**: 78-81.
- Gadakh, S.S., Dethe, A.M., Kathale, M.N. and Kahate N.S. 2013. Genetic diversity for yield and its component traits in green gram [*Vigna radiata* (L.) Wilczek]. *Journal of Crop and Weed*, **9**(1): 106-10.
- Idress, A., Sadiq, M. S., Hanif, M., Abbas, G. and Haider, S. 2006. Genetic parameters and path coefficient analysis in mutated generation of mungbean (*Vigna radiata* L. Wilckez.). *Journal of Agricultural Research*, **44**: 181-191.
- Jeberson, M.S., Shashidhar, K.S. and Iyanar, K. 2015. Estimation of genetic parameters for phenological traits in mungbean. *Trends in Biosciences*, **8**(21): 5741-5745.
- Johnson, H.F., Robinson, H.F. and Comstock, R.E. 1955. Estimates of genetic and environmental variability in soybean. *Agronomy Journal*, **47**: 314-318.



- Karpechenko, G.D. 1925. On the chromosomes of Phaseolinae. *Bull. Appl. Bot. Genet. Breed.* (Leningrad), **14**: 143–148.
- Kumar, B.S., Prakash, M. and Gokul Krishna, J. 2010. Combining ability analysis in mungbean (*Vigna radiata* (L.) Wilczek). *Crop Improvement*, **37**(2): 165-167.
- Kumar, R. Tripathi, I.D., Luthra, O.P. and Kumar, R. 1996. Estimates of variability parameters in wheat. *Haryana Agric. Univ. J. Res.*, **26**: 223-226.
- Lakshminarayanareddy, V., Reddisekhar, M., Raja Reddy, K. and Hariprasad Reddy, K. 2003. Genetic variability for yield and its components in mungbean (*Vigna radiata* (L.) Wilczek). *Legume Research*, **26**(4): 300-302.
- Makeen, K., Abraham, G., Jan, A. and Singh, A.K. 2007. Genetic variability and correlation studies on yield and its components in mungbean (*Vigna radiata* (L.) Wilczek). *Journal of Agronomy*, **6**(1): 216 - 218. [[Cross Ref](#)]
- Mehandi, S., Mishra, S.P., Tripathi, R.C. and Singh, I.P. 2018. Genetic variability, heritability and genetic advance for yield and its related traits in mungbean [*Vigna radiata* (L.) Wilczek] genotypes. *International Journal of Current Microbiology and Applied Sciences*, 3818-3824.
- Nand, M.J. and Anuradha, C. 2013. Genetic variability, correlation and path coefficient analysis for yield and yield components in mung bean [*Vigna radiata* (L.) Wilczek]. *Journal of Research, ANGRAU*, **41**(3): 31-39.
- Narasimhulu, R., Naidu, N. V., Priya, M. S., Rajarajeswari, V. and Reddy, K.H.P. 2013. Genetic variability and association studies for yield attributes in mungbean (*Vigna radiata*). *Indian Journal of Plant Sciences*, **2**(3): 82-86.
- Panse, V.G. and Sukhatme, P.V. 1978. *Statistical Methods for Agricultural Workers*, ICAR, New Delhi.
- Pavan, K., Reddy, P. and Mehta, C.M. 2019. Estimation of variability through genetic parameters and identification of superior pure lines for yield attributing traits in green gram [*Vigna radiata* (L.) Wilczek]. *Journal of Pharmacognosy and Phytochemistry*, SP3: 55-61.
- Pawar, I.S., Redhu, A.S., Singh, I., Yunus, M. and Singh, I. 2001. Effectiveness of selection procedure in wheat. *Haryana Agril. Univ. J. Res.*, **31**: 119-121.
- Pawar, S.D., Thete, R.Y. and Dumbre, A.D. 1988. Estimates of genetic variability parameters in F2 population of wheat. *J. Maharashtra Agric. Univ.*, **13**: 210-211.
- Prajapati, S.S., Singh, S.K., Shrivastava, M.K., Biswal, M., Kumar, P., Rahangdale, S. and Behera, K. 2023. Evaluation of genetic parameters for yield and yield related traits in mungbean [*Vigna radiata* (L.) Wilczek]. *The Pharma Innovation Journal*, **12**(3): 982-986.
- Ramachandra, P. and Lavanya, G.R. 2017. Variability, heritability genetic advance and correlation coefficients for yield component characters and seed yield in Green gram (*Vigna radiata* (L.) Wilczek). *Journal of Pharmacognosy and Phytochemistry*, **6**(4): 1202-1205.
- Robinson, H.F., Comstock, R.E. and Harvey, P.H. 1949. Estimates of heritability and the degree of dominance in corn. *Agron. J.*, **41**: 353-359. [[Cross Ref](#)]
- Sirohi, S.P.S., Tyagi, M., Sirohi, S. and Mohit. 2018. Assessment of genetic variability and character association in promising genotypes of mungbean [*Vigna radiata* (L.) Wilczek]. *Journal of Pharmacognosy and Phytochemistry*, **7**(5): 2513-2516.
- Sofia, S., Reddy, D.M., ShantiPriya, M. and Latha, P. 2017. Genetic variability studies on physiological and yield related traits in mungbean [*Vigna radiata* (L.) Wilczek]. *Bulletin of Environment, Pharmacology and Life Sciences*, **6**(2): 252-255.
- Susmitha, D. and Jayamani, P. 2018. Genetic variability studies for yield and its contributing traits in green gram [*Vigna radiata* (L.) Wilczek]. *Electronic Journal of Plant Breeding*, **9**(20): 716-722. [[Cross Ref](#)]
- Tabasum, A., Saleem, M. and Aziz, I. 2010. Genetic variability, trait association and path analysis of yield and yield components in mung bean [*Vigna radiata* (L.) Wilczek]. *Pakistan Journal of Botany*, **42**(6): 3915-3924.
- Tiwari, A., Mishra, S.P. and Nag, S.K. 2014. Genetic variability, heritability and genetic advance studies for yield and its components in mungbean [*Vigna radiata* (L.) Wilczek]. *Trends in Biosciences*, **7**(1): 944-946.
- Vavilov, N.I. 1926. Origin variation, immunity of cultivated plants. *Chronica Bot*, **13**: 364.
- Yadav, R., Ram, B.L., Suresh, B.G. and Lavanya, G.R. 2017. Studies on genetic variability, character association for yield and yield components in green gram [*Vigna radiata* (L.) Wilczek]. *Journal of Pharmacognosy and Phytochemistry*, **6**(4): 1988-1991.
- Yoseph, T., Mekbib, F., Fenta, B.A. and Tadele, Z. 2022. Genetic variability, heritability, and genetic advance in mung bean [*Vigna radiata* (L.) Wilczek] genotypes. *Ethiopian Journal of Crop Science*, **9**(2).